

# Particulate Matter 10 (PM10): Persistence and Trends in Eight European Capitals

*Guglielmo Maria Caporale, Luis A. Gil-Alana, Nieves Carmona-González*

## **Impressum:**

CESifo Working Papers

ISSN 2364-1428 (electronic version)

Publisher and distributor: Munich Society for the Promotion of Economic Research - CESifo GmbH

The international platform of Ludwigs-Maximilians University's Center for Economic Studies and the ifo Institute

Poschingerstr. 5, 81679 Munich, Germany

Telephone +49 (0)89 2180-2740, Telefax +49 (0)89 2180-17845, email [office@cesifo.de](mailto:office@cesifo.de)

Editor: Clemens Fuest

<https://www.cesifo.org/en/wp>

An electronic version of the paper may be downloaded

- from the SSRN website: [www.SSRN.com](http://www.SSRN.com)
- from the RePEc website: [www.RePEc.org](http://www.RePEc.org)
- from the CESifo website: <https://www.cesifo.org/en/wp>

# Particulate Matter 10 (PM10): Persistence and Trends in Eight European Capitals

## Abstract

This paper examines the statistical properties of daily PM10 in eight European capitals (Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris) over the period 2014-2020 by applying a fractional integration framework; this is more general than the standard approach based on the classical dichotomy between I(0) stationary and I(1) non-stationary series used in most other studies on air pollutants. All series are found to be characterised by long memory and fractional integration, with orders of integration in the range (0, 1), which implies that mean reversion occurs and shocks do not have permanent effects. Persistence is highest in the case of Brussels, Amsterdam and London. The presence of negative trends in Brussels, Paris and Berlin indicates some degree of success in reducing pollution in these capitals.

JEL-Codes: C220, Q530.

Keywords: fractional integration, long memory, persistence, trends, air pollutants, PM<sub>10</sub>.

*Guglielmo Maria Caporale\**  
*Department of Economics and Finance*  
*Brunel University London*  
*United Kingdom – London, UB8 3PH*  
*Guglielmo-Maria.Caporale@brunel.ac.uk*

*Luis A. Gil-Alana*  
*University of Navarra*  
*Pamplona / Spain*  
*alana@unav.es*

*Nieves Carmona-González*  
*Universidad Francisco de Vitoria*  
*Madrid / Spain*

\*corresponding author

June 2020

Prof. Luis A. Gil-Alana gratefully acknowledges financial support from the MINEIC-AEI-FEDER ECO2017-85503-R project from ‘Ministerio de Economía, Industria y Competitividad’ (MINEIC), ‘Agencia Estatal de Investigación’ (AEI) Spain and ‘Fondo Europeo de Desarrollo Regional’ (FEDER). He also acknowledges financial support from internal Projects of the Universidad Francisco de Vitoria.

## 1. Introduction

Particulate matter (PM) are microscopic particles of solid or liquid matter suspended in the air. Its sources can be natural or anthropogenic. It has a significant impact on both climate and precipitation, and therefore has both health and social costs. In particular, it affects the amount of incoming solar radiation and outgoing terrestrial radiation. The coarse particles can have a diameter between 2.5 and 10 micrometers ( $\mu\text{m}$ ) ( $\text{PM}_{10}$ ) and are known to be a very harmful form of air pollution given their ability to penetrate into the lungs and blood streams and cause respiratory and heart diseases as well as premature death. Various countries have therefore set limits for particulars in the air, which are emitted during the combustion of vehicle engine fuels, braking and tyre wearing. In particular, the European Union has defined in a series of directives the acceptable limits for exhaust emissions of new vehicles sold in the European Union and EEA member states.

Numerous studies have analysed the connection between pollution and harmful health effects (e.g., Schwartz and Marcus, 1990; Anderson et al., 1996; Atkinson et al., 1999; Gardner and Dorling, 1999). The present study contributes to another branch of the literature which focuses instead on modelling various pollutants such as sulphur dioxide ( $\text{SO}_2$ ), nitrogen dioxide ( $\text{NO}_2$ ), carbon monoxide (CO), ozone ( $\text{O}_3$ ),  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$ . For instance, Zamri et al. (2009) applied the Box-Jenkins ARIMA approach to model CO and  $\text{NO}_2$  in Malaysia and found an upward trend. Li et al. (2017) analysed air quality in Beijing from 2014 to 2016 using the Spatio-temporal Deep Learning (StDL) model, the Time Delay Neural Network (TDNN) model, the ARMA model, the Support Vector Regression (SVR) model, and the Long Short-Term Memory Neural Network Extended (LSTME) model, and concluded that the LSTME model is the most suitable one for time series characterised by long-term dependence with optimal time delays.

Naveen and Anu (2017) studied air quality in India using ARIMA, seasonal ARIMA (SARIMA) and other models. Pan and Chen (2008) is one of the few studies using long-memory AutoRegressive Fractional Integrated Moving Average (ARFIMA) models for air pollution data (in Taiwan) and concluding that these are more accurate than AutoRegressive Integrated Moving Average (ARIMA) models.

It is clearly important to investigate the dynamics of air pollution to develop suitable models for prediction purposes and design policies to manage air quality. This paper examines the statistical properties of daily  $PM_{10}$  in eight European capitals (Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris) over the period 2014-2020 by applying a fractional integration framework that is more general than the standard approach based on the classical dichotomy between  $I(0)$  stationary and  $I(1)$  non-stationary series used in the vast majority of previous studies on air pollutants, since it allows for fractional as well as integer degrees of differentiation and thus for a much wider set of stochastic behaviours. In particular, it enables the researcher to analyse the long-memory properties of the series of interest and the possible presence of trends, to test for mean reversion, and to measure the degree of persistence and the speed of adjustment to the long-run equilibrium level. Therefore, it provides information about whether the effects of shocks are transitory or permanent, which is a crucial piece of information for adopting appropriate policy measures.

The remainder of the paper is structured as follows: Section 2 outlines the methodology used for the analysis; Section 3 describes the data; Section 4 presents the empirical results; Section 5 offers some concluding remarks.

## 2. Methodology

As mentioned above, we adopt a long-memory approach based on fractional integration. Long memory is a feature of time series that are characterised by a high degree of dependence between observations which are far apart in time. It has been found to be displayed by many time series in different fields such as climatology (Gil-Alana, 2005; 2008; 2017; Vyushin and Kushner, 2009; Franzke, 2012; Ludescher et al., 2016; Bunde, 2017; Yuan et al., 2019); environmental sciences (Barros et al., 2016; Gil-Alana et al., 2016; Tiwari et al., 2016; Gil-Alana and Solarin, 2018); economics and finance (Gil-Alana and Moreno, 2012; Abritti et al., 2017); etc.

There exist a variety of statistical models that can describe this type of behaviour; a very popular one among time series analysts is based on the concept of fractional integration, which occurs when the number of differences required to make a series stationary  $I(0)$  is a fractional value. More precisely, a time series is said to be integrated of order  $d$  or  $I(d)$  if it can be expressed as:

$$(1 - B)x_t = u_t, \quad t = 1, 2, \dots, \quad (1)$$

where  $B$  is the backshift operator ( $Bx_t = x_{t-1}$ ), the differencing parameter  $d$  can be any real value, and  $u_t$  is  $I(0)$  defined as a covariance stationary process with a spectral density function that is positive and bounded at all frequencies in the spectrum. This framework encompasses different cases such as short memory ( $d = 0$ ), stationary long memory ( $0 < d < 0.5$ ), non-stationary though mean-reverting processes ( $0.5 \leq d < 1$ ), unit roots ( $d = 1$ ) and explosive patterns ( $d \geq 1$ ).

## 3. Data

The series analysed is the daily average air quality taken from the World Air Quality Index (WAQI) at <https://aqicn.org/map/world/es/>. All data have been converted using

the US EPA standard (United States Environmental Protection Agency). Specifically, we use daily data for the past 7 years (2014-2020) concerning eight European capitals: Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris. The series represents the daily level of air quality (PM10) measured in micrograms per cubic meter of air ( $\mu\text{g}/\text{m}^3$ ). The WAQI data come from the following original sources: Madrid: <http://www.mambiente.madrid.es/opencms/opencms/calibre/> (Ayuntamiento de Madrid); Paris: <http://www.airparif.asso.fr/> (AirParif - Association de surveillance de la qualité de l'air en Île-de-France); Amsterdam: <https://www.luchtmeetnet.nl/> (RIVM); Luxembourg: <https://environnement.public.lu/fr.html> (Portail de l'Environnement du Grand-duché de Luxembourg); London: <https://uk-air.defra.gov.uk/> (UK-AIR, air quality information resource - Defra, UK); Helsinki: <https://www.ilmatieteenlaitos.fi/ilmanlaatu> (Ilmanlaatu Suomessa); Brussels: <https://www.irceline.be/en/> (Belgian Interregional Environment Agency); Berlin: <https://www.berlin.de/senuvk/umwelt/luftqualitaet/> (Luftqualität).

**[TABLE 1 ABOUT HERE]**

Table 1 reports the sample periods for each capital and provides some descriptive statistics for each series. It can be seen that Paris exhibits the highest mean value, while Helsinki has the lowest. Paris also has the most volatile series, whilst Luxembourg has the least volatile.

**4. Empirical Results**

We estimate the following model:

$$y_t = \alpha + \beta t + x_t, \quad (1 - B)^d x_t = u_t, \quad t = 1, 2, \dots, \quad (2)$$

where  $y_t$  stands for  $PM_{10}$  in each European capital in turn;  $B$  is the backshift operator, and  $x_t$  is an  $I(d)$  process such that the error term  $u_t$  is  $I(0)$ ; the disturbances are assumed to follow a white noise (see Tables 2 and 3) and an autocorrelated process (see Tables 4 and 5) in turn, where the latter is modelled using the exponential spectral framework of Bloomfield (1973). In all cases we display the estimated values of  $d$  (and their associated 95% confidence bands) for three different specifications: (i) no deterministic terms in (1), i.e., we impose the restriction  $\alpha = \beta = 0$  (the results for this case are reported in the second column in Tables 2 and 4); (ii) an intercept only (see the third column in both tables); (iii) an intercept and a linear time trend (see the fourth column in both tables). The estimated values reported in bold in these tables are those corresponding to our preferred specification, which has been selected on the basis of the statistical significance of the regressors.

**[TABLES 2 AND 3 ABOUT HERE]**

When assuming that  $u_t$  is a white noise, the intercept is found to be the only significant deterministic term in all cases; the estimated values of  $d$  are in the interval  $(0, 1)$ , which implies long memory and fractional integration. They range between 0.39 (Amsterdam) and 0.62 (Madrid). For Amsterdam, the values are all within the stationary range ( $d < 0.5$ ); for Brussels, London, Paris and London, they are around the stationary boundary ( $d = 0.5$ ), while non-stationarity ( $d \geq 0.5$ ) is found in the case of Helsinki, Berlin and Madrid.

**[TABLES 4 AND 5 ABOUT HERE]**



When allowing for autocorrelation (Tables 4 and 5) the time trend appears to be negative and statistically significant in the case of Brussels, Berlin and Paris, which might reflect the anti-pollution policies adopted in these three capitals. In particular, a low emission zone (LEZ) was established in the Brussels region with the aim of meeting the European air quality standards and emission ceilings; in Berlin the German Climate Action Plan 2050 is being implemented to control air pollution by laying down environmental quality standards and emission reduction requirements; a LEZ based on Euro norm vehicle classification has also been introduced in Paris.

The estimated values of  $d$  are once more in the interval  $(0, 1)$ , though they are now significantly smaller than in the previous case. In fact, they are all within the stationary range, specifically between 0.22 (Luxembourg) and 0.33 (Helsinki and Paris). These lower estimates are likely to reflect the competition between the fractional integration and Bloomfield parameters in describing time dependence between the observations. Both sets of estimates, under the assumption of white noise and autocorrelated errors respectively, indicate that the degree of persistence is highest in the case of Brussels, Amsterdam and London, and lowest in the case of Helsinki, Berlin, and Madrid; thus, the effects of shocks are more long-lived in the former capitals.

## **5. Conclusions**

This paper has used fractional integration methods to obtain evidence on persistence and time trends in  $PM_{10}$  in eight European capitals (Amsterdam, Berlin, Brussels, Helsinki, London, Luxembourg, Madrid and Paris). This approach is more general than the standard ones used in most of the literature on air pollutants and thus is more informative about the time series properties of the series of interest. The results indicate that all of them display fractional integration with orders of integration in the range

(0,1); this implies that mean reversion occurs and shocks do not have permanent effects. However, the degree of persistence is different in the eight capitals examined; in particular, the effects of shocks take longer to die away in the case of Brussels, Amsterdam and London. Such evidence should be taken into account by policy makers aiming to design effective measures to reduce pollution.

The estimated values of  $d$  are lower under the assumption of autocorrelated errors; in this case three of the capitals examined (Brussels, Paris and Berlin) exhibit statistically significant negative time trends, which suggests that the policies they have adopted to reduce pollution (such as the establishment of LEZs) have been successful, at least to some extent.

Other statistical properties of  $PM_{10}$  that could be investigated are seasonality, non-linearities and structural breaks; the forecasting properties of rival models could also be examined; all these issues are left for future work.

## References

- Abritti, M., L.A. Gil-Alana, Y. Lovcha and A. Moreno (2017) Term structure persistence, *Journal of Financial Econometrics* 14, 331-352.
- Anderson HR, Ponce de Leon A, Martin-Bland J, Bower J.S., Strachan DP (1996) Air pollution and daily mortality in London: 1987-92, *BMJ* 312:665-669.  
<http://doi:10.1136/bmj.312.7032.665>
- Atkinson RW, Anderson HR, Strachan DP, Bland JM, Bremner SA, Ponce de Leon A (1999) Short-term associations between outdoor air pollution and visits to accident and emergency departments in London for respiratory complaints *Eur Respir J* 13(2):257–265. [http://doi: 10.1183/09031936.99.1322579](http://doi:10.1183/09031936.99.1322579)
- Barros, C.P., L.A. Gil-Alana and F. Perez de Gracia (2016), Stationarity and long range dependence of carbon dioxide emissions. Evidence from disaggregated data, *Environmental and Resources Economics* 63, 1, 45-56.
- Bunde, A. (2017), Long-term memory in climate: Detection, extreme events and significance of trends, Chapter 11 in *Nonlinear and Stochastic Climate Dynamics*, edited by Christian L.E. Franzke and Terence O’Kane, Cambridge University Press
- Franzke, C. (2012), Nonlinear trends, long-range dependence, and climate noise properties of surface temperature, *Journal of Climate*, 25(12), 4172–4183.
- Gardner MW, Dorling SR (1999) Neural network modelling and prediction of hourly NO<sub>x</sub> and NO<sub>2</sub> concentrations in urban air in London. *Atmos Environ* 33:709-719.  
[https://doi.org/10.1016/S1352-2310\(98\)00230-1](https://doi.org/10.1016/S1352-2310(98)00230-1)
- Gil-Alana, L.A. (2005), Statistical modeling of the temperatures in the Northern hemisphere using fractional integration techniques, *Journal of Climate* 27, 10, 3477-3491.
- Gil-Alana, L.A. (2008), Time trend estimation with breaks in temperature series, *Climatic Change* 89, 3, 325-337.
- Gil-Alana, L.A., (2017) Alternative modeling approaches for the ENSO time series. Persistence and seasonality, *International Journal of Climatology* 37, 5, 2354-2363.
- Gil-Alana, L.A., R. Gupta and F. Perez de Gracia (2016), Modelling persistence of carbon emissions allowance prices, *Renewable and Sustainable Energy Review* 55, 221-226.
- Gil-Alana, L.A. and A. Moreno (2012), Uncovering the US term premium. An alternative route, *Journal of Banking and Finance* 36, 4, 1181-1193.
- Gil-Alana, L. A., and Solarin, S. A. (2018). Have US environmental policies been effective in the reduction of US emissions? A new approach using fractional integration. *Atmospheric Pollution Research*, 9(1), 53-60.

- Li X, Peng L, Yao X, Cui S, Hi Y, You C, Chi T (2017). Long short-term memory neural network for air pollutant concentration predictions: Method development and evaluation. *Environ Pollut* 231:1997-1004. <https://doi.org/10.1016/j.envpol.2017.08.114>
- Ludescher, J., A. Bunde, C.L. Franzke and H. J. Schellnhuber (2016). Long-term persistence enhances uncertainty about anthropogenic warming of Antarctica, *Climate Dynamics*, 46(1–2), 263–271, doi: 10.1007/s00382-015-2582-5
- Naveen V, Anu N (2017) Time series analysis to forecast air quality indices in Thiruvananthapuram District, Kerala, India. *J Eng Res Appl* 7(6): 66-84. <https://doi: 10.9790/9622-0706036684>
- Pan JN, Chen ST (2008) Monitoring long-memory air quality data using ARFIMA model, *Environmetrics* 19 (2): 209–219. <https://doi.org/10.1002/env.88>
- Schwartz O, Marcus A (1990) Mortality and air pollution in London: A time series analysis. *Am J Epidemiol* 131(1):185–194. <https://doi: 10.1136/jech.54.10.750>
- Tiwari, A.K., Kyophilavong, P. and Albulescu, C.T., (2016), Testing the stationarity of CO<sub>2</sub> emissions series in Sub-Saharan African countries by incorporating nonlinearity and smooth breaks, *Research in International Business and Finance*, 37, 527-540.
- Vyushin, D. I. and P.J. Kushner (2009), Power-law and long-memory characteristics of the atmospheric general circulation, *Journal of Climate*, 22(11), 2890–2904.
- Yuan, N., Y. Huang, J. Duan, C. Zhu, E. Xoplaki and J. Luterbacher (2019), On climate prediction; how much can we expect from climate memory?, *Climate Dynamics* 52, 1-2, 855-864.
- Zamri IM, Roziah Z, Marzuki I, Muhd SL (2009) Forecasting and Time Series Analysis of Air Pollutants in Several Area of Malaysia. *Am J Environ Sci* 5:625-632. <https://doi:10.3844/ajessp.2009.625.632>

**Table 1: Descriptive statistics**

<b>AMSTERDAM</b>		<b>BERLIN</b>	
Start date	End date	Start date	End date
01/01/2014	13/06/2020	20/08/2014	13/06/2020
<i>No. of observations:</i> 2145 <i>Mean:</i> 25.7 <i>Standard deviation:</i> 11.1 <i>Variance:</i> 124.1 <i>Min./Max.:</i> 8 / 263 <i>Range:</i> 255		<i>No. of observations:</i> 2114 <i>Mean:</i> 28.0 <i>Standard deviation:</i> 11.9 <i>Variance:</i> 140.6 <i>Min./Max.:</i> 8 / 95 <i>Range:</i> 87	
<b>BRUSSELS</b>		<b>HELSINKI</b>	
Start date	End date	Start date	End date
31/12/2013	13/06/2020	02/05/2014	13/06/2020
<i>No. of observations:</i> 2334 <i>Mean:</i> 24.0 <i>Standard deviation:</i> 11.8 <i>Variance:</i> 140.4 <i>Min./Max.:</i> 1 / 100 <i>Range:</i> 99		<i>No. of observations:</i> 2214 <i>Mean:</i> 18.1 <i>Standard deviation:</i> 10.0 <i>Variance:</i> 99.9 <i>Min./Max.:</i> 3 / 90 <i>Range:</i> 87	
<b>LONDON</b>		<b>LUXEMBOURG</b>	
Start date	End date	Start date	End date
31/12/2013	13/06/2020	19/06/2015	13/06/2020
<i>No. of observations:</i> 2353 <i>Mean:</i> 26.5 <i>Standard deviation:</i> 9.9 <i>Variance:</i> 97.1 <i>Min./Max.:</i> 5 / 89 <i>Range:</i> 84		<i>No. of observations:</i> 1583 <i>Mean:</i> 19.5 <i>Standard deviation:</i> 6.8 <i>Variance:</i> 46.2 <i>Min./Max.:</i> 2 / 52 <i>Range:</i> 50	
<b>MADRID</b>		<b>PARIS</b>	
Start date	End date	Start date	End date
31/12/2013	13/06/2020	31/12/2013	13/06/2020
<i>No. of observations:</i> 2323 <i>Mean:</i> 24.3 <i>Standard deviation:</i> 11.7 <i>Variance:</i> 138.0 <i>Min./Max.:</i> 5 / 160 <i>Range:</i> 155		<i>No. of observations:</i> 2227 <i>Mean:</i> 39.3 <i>Standard deviation:</i> 14.5 <i>Variance:</i> 210.5 <i>Min./Max.:</i> 6 / 122 <i>Range:</i> 116	

**Table 2: Estimates of d: White noise errors**

Series	No deterministic terms	An intercept	An intercept and a time trend
AMSTERDAM	0.42 (0.39, 0.46)	<b>0.39 (0.35, 0.44)</b>	0.39 (0.35, 0.44)
BERLIN	0.63 (0.58, 0.68)	<b>0.61 (0.56, 0.67)</b>	0.61 (0.56, 0.67)
BRUSSELS	0.52 (0.48, 0.56)	<b>0.50 (0.46, 0.55)</b>	0.50 (0.46, 0.55)
HELSINKI	0.57 (0.52, 0.61)	<b>0.54 (0.50, 0.59)</b>	0.54 (0.50, 0.59)
LONDON	0.54 (0.50, 0.59)	<b>0.52 (0.47, 0.57)</b>	0.52 (0.47, 0.57)
LUXEMBOURG	0.56 (0.51, 0.62)	<b>0.54 (0.48, 0.61)</b>	0.54 (0.48, 0.61)
MADRID	0.63 (0.58, 0.68)	<b>0.62 (0.57, 0.67)</b>	0.62 (0.57, 0.67)
PARIS	0.55 (0.51, 0.59)	<b>0.53 (0.49, 0.58)</b>	0.53 (0.49, 0.58)

We report the estimates of d and its 95% confidence band (in parenthesis). In bold, the selected specification for each series.

**Table 3: Estimated coefficients in the selected model: White noise errors**

Series	d (95% band)	Intercept (t-value)	Time trend
AMSTERDAM	0.39 (0.35, 0.44)	26.0450 (8.77)	---
BERLIN	0.61 (0.56, 0.67)	25.3864 (3.93)	---
BRUSSELS	0.50 (0.46, 0.55)	20.9146 (4.17)	---
HELSINKI	0.54 (0.50, 0.59)	20.4913 (4.57)	---
LONDON	0.52 (0.47, 0.57)	23.0696 (5.23)	---
LUXEMBOURG	0.54 (0.48, 0.61)	15.6394 (4.61)	---
MADRID	0.62 (0.57, 0.67)	16.5249 (2.59)	---
PARIS	0.53 (0.49, 0.58)	32.9226 (5.16)	---

The values in parenthesis in column 3 are the corresponding t-values.

**Table 4: Estimates of d: Autocorrelated errors**

Series	No deterministic terms	An intercept	An intercept and a time trend
AMSTERDAM	0.34 (0.29, 0.39)	<b>0.26 (0.21, 0.30)</b>	0.25 (0.20, 0.30)
BERLIN	0.38 (0.33, 0.43)	0.30 (0.25, 0.36)	<b>0.29 (0.23, 0.36)</b>
BRUSSELS	0.33 (0.28, 0.38)	0.26 (0.21, 0.32)	<b>0.25 (0.21, 0.31)</b>
HELSINKI	0.40 (0.35, 0.45)	<b>0.33 (0.29, 0.39)</b>	0.33 (0.29, 0.38)
LONDON	0.34 (0.30, 0.39)	<b>0.26 (0.21, 0.31)</b>	0.27 (0.22, 0.30)
LUXEMBOURG	0.32 (0.28, 0.38)	<b>0.22 (0.16, 0.29)</b>	0.21 (0.15, 0.28)
MADRID	0.34 (0.30, 0.39)	<b>0.30 (0.25, 0.35)</b>	0.30 (0.25, 0.35)
PARIS	0.41 (0.37, 0.46)	0.33 (0.28, 0.38)	<b>0.33 (0.28, 0.38)</b>

We report the estimates of d and its 95% confidence band (in parenthesis). In bold, the selected specification for each series.

**Table 5: Estimated coefficients in the selected model: Autocorrelated errors**

Series	d (95% band)	Intercept (t-value)	Time trend (t-value)
AMSTERDAM	0.26 (0.21, 0.30)	25.6832 (21.18)	---
BERLIN	0.29 (0.23, 0.36)	32.6088 (16.38)	-0.0046 (-2.92)
BRUSSELS	0.25 (0.21, 0.31)	26.3621 (15.22)	-0.0022 (1.97)
HELSINKI	0.33 (0.29, 0.39)	19.6888 (12.88)	---
LONDON	0.26 (0.21, 0.31)	25.9193 (24.21)	---
LUXEMBOURG	0.22 (0.16, 0.29)	19.2309 (34.57)	---
MADRID	0.30 (0.25, 0.35)	23.5407 (16.78)	---
PARIS	0.33 (0.28, 0.38)	42.2180 (13.79)	-0.0044 (-1.98)

The values in parenthesis in columns 3 and 4 are the corresponding t-values for the intercept and the time trend respectively.