

Endogenous Risk Attitudes

Nick Netzer, Arthur Robson, Jakub Steiner, Pavel Kocourek

Impressum:

CESifo Working Papers

ISSN 2364-1428 (electronic version)

Publisher and distributor: Munich Society for the Promotion of Economic Research - CESifo GmbH

The international platform of Ludwigs-Maximilians University's Center for Economic Studies and the ifo Institute

Poschingerstr. 5, 81679 Munich, Germany

Telephone +49 (0)89 2180-2740, Telefax +49 (0)89 2180-17845, email office@cesifo.de

Editor: Clemens Fuest

<https://www.cesifo.org/en/wp>

An electronic version of the paper may be downloaded

- from the SSRN website: www.SSRN.com
- from the RePEc website: www.RePEc.org
- from the CESifo website: <https://www.cesifo.org/en/wp>

Endogenous Risk Attitudes

Abstract

In a model inspired by neuroscience, we show that constrained optimal perception encodes lottery rewards using an S-shaped encoding function and over-samples low-probability events. The implications of this perception strategy for behavior depend on the decision-maker's understanding of the risk. The strategy does not distort choice in the limit as perception frictions vanish when the decision-maker fully understands the decision problem. If, however, the decision-maker underrates the complexity of the decision problem, then risk attitudes reflect properties of the perception strategy even for vanishing perception frictions. The model explains adaptive risk attitudes and probability weighting as in prospect theory and, additionally, predicts that risk attitudes are strengthened by time pressure and attenuated by anticipation of large risks.

JEL-Codes: D810, D870, D910.

Keywords: endogenous preferences, probability distortions, misspecified learning.

Nick Netzer
University of Zurich / Switzerland
nick.netzer@econ.uzh.ch

Arthur Robson
Simon Fraser University
Burnaby / British Columbia / Canada
robson@sfu.ca

Jakub Steiner
University of Zurich / Switzerland
jakubsteiner77@gmail.com

Pavel Kocourek
Cerge-Ei
Prague / Czech Republic
pakocica@gmail.com

January 25, 2022

We have benefited from comments of Cary Frydman, Ryan Webb and various seminar audiences. Giorgi Chavchanidze and Pavel Ilinov provided excellent research assistantship. This work was supported by the Alfred P. Sloan Foundation and the NOMIS Foundation (Netzer), by SSHRC grant 435-2013-0426 (Robson) and by ERC grant 770652 (Steiner).

1 Introduction

Although economists usually take preferences as exogenous and fixed, there is compelling evidence that these change with the context. For choices over gambles, we know at least since Kahneman and Tversky (1979) that risk attitudes are not fixed: the steep part of the S-shaped utility function in prospect theory adapts to the status quo. Rabin’s (2000) paradox provides another challenge for stable risk attitudes: choices over small and large risks are best represented by distinct Bernoulli utility functions. Risk attitudes are further modulated by external factors such as time pressure or framing (e.g., Kahneman, 2011). An additional, well-known anomaly involves the overweighting of small objective probability events relative to more likely events (Kahneman and Tversky, 1979).

In this paper, we explain these anomalies in a unifying way. In our model, endogenous risk attitudes and probability weighting are the consequence of constrained optimal perception of lotteries, combined with a possible misspecification of the structure of the risk.

Our procedural choice model is inspired by the literature on optimal coding from neuroscience. A risk-neutral decision-maker (DM) chooses between a lottery and a safe option. For illustration, consider the vivid example from Savage (1954) in which choosing the lottery represents purchase of a convertible car, the enjoyment of which depends on the random weather conditions. The DM knows the probabilities of the possible states of the world (the different weather conditions), observes the value of the safe option (the price of the car), but faces a friction in processing the rewards of the lottery in the different states (the weather-dependent enjoyments). She learns about the reward vector by sampling signals. Each signal is a reward of the lottery in a respective state encoded via a non-linear encoding function that maps rewards to their mental representations (e.g. neural firing rates), then perturbed by additive Gaussian noise. Noisy mental representations is what the DM (or her brain) collects when extracting information from the description of the lottery, when retrieving experiences from memory, or when processing the experiences provided by others (e.g. the car dealer). After she has observed a collection of signals, the DM forms a maximum-likelihood or Bayesian estimate of the value of the lottery and takes the optimal decision.

The DM optimizes the perception strategy – the encoding function and the sampling frequencies of all states – for a given distribution of decision problems. Choice of the perception strategy is a specific form of an attention allocation problem. Our DM is akin to an engineer who measures a physical input by reading off the position of a needle on a meter. The engineer can choose the measurement function that maps the physical input to the needle position. If the needle position has a stochastic component, then the engineer can increase the precision of her measurement for a specific range of inputs by making the

measurement function steep in this range. Our DM can increase the precision of her reward perception for a specific range of rewards by making the encoding function steep in this range. Since the range of possible mental representations is finite, the encoding function cannot be steep everywhere. Further, our DM can allocate attention to a specific state of the world by sampling it frequently, but this comes at the cost of sampling other states less frequently.

The model explains adaptive S-shaped encoding of rewards and over-sampling of low-probability states as jointly optimal. The implications of the perception strategy for behavior are subtle. As the perception data become rich and approximate full information, behavior becomes risk-neutral whenever the DM understands the structure of the risk she faces, and hence learns about it in a correctly specified model. However, the perception strategy induces non-trivial risk attitudes when the DM applies a simplifying model to the encountered risk. In that case, the encoding function takes on the role of a Bernoulli utility function and over-sampling of small probability states translates into overweighting of small probabilities, generating risk attitudes akin to those from prospect theory.

We analyze the limit of rich perception data, motivated by two considerations. First, while human perception is inherently noisy, imprecisions can be partially mitigated by collecting more data, in particular when the stakes are large. Our limiting results are a useful approximation when noise is small relative to the stakes of the decision problem. Second, the limit is tractable. We prove that the expected loss from misperception, relative to choice under complete information, is approximately the mean squared error in perception of the lottery value, integrated over all decision problems in which the lottery value ties with the safe option. The conditioning on ties arises endogenously. Accuracy of perception has instrumental value for choice, and choice is trivial except where the values of two options are nearly equal, given that information is nearly complete.

We then derive the perception strategy for which the mean squared error over ties goes to zero most quickly as perception data become rich. This problem has a unique solution which is characterized by intuitive optimality conditions. For the plausible case of unimodal symmetric reward densities, we show that an S-shaped encoding function and over-sampling of low-probability states are jointly optimal. The DM chooses the encoding function to be steep near the modal rewards and flatter towards the tails of the reward distribution. She thus perceives the reward values typical for her environment relatively precisely, at the expense of precision at the tail rewards. Conditioning on ties induces a statistical association between tail rewards and low-probability states, because tail rewards in high-probability states typically result in very attractive or unattractive lotteries rather than in ties. Thus, tail rewards arise relatively often in the low-probability states at ties. Hence, the DM

with an S-shaped encoding function relatively often struggles to estimate the rewards from low-probability states. It is optimal to compensate for this by over-sampling such states. To illustrate, consider the decision whether to take a flight. The DM may struggle to comprehend a possible aviation accident since her apprehension is not well adapted to such extreme events. Such extreme events have only small probabilities in lotteries involved in non-trivial comparisons – otherwise choice would be trivial. Paying disproportionate attention to the contingency of the aviation accident then optimally compensates for the struggle of its apprehension.

We then turn to the behavioral consequences of the perception strategy. If the DM learns in a correctly specified model, then the perception strategy affects precision of perception rather than risk attitudes. For instance, a locally steep encoding function translates into locally precise decoding of the reward values rather than into high marginal utility. Similarly, oversampling of a state translates for a well-specified DM into increased precision of the state’s perceived reward rather than in an increased subjective probability of the state. We thus obtain the prediction that the choices of a well-specified DM become undistorted in the limit as the number of signals grows. For example, the decisions of a financial expert with access to rich data in her domain of expertise will not be influenced by her perception strategy.

On the other hand, if the DM interprets the perception data in a naturally misspecified model, then a tight connection between the perception strategy and risk attitudes arises. To illustrate the main idea, consider again the engineer who observes the needle position on her meter and knows that the position is a non-linear function of the measured input. But now assume that the needle trembles due to stochasticity of the input. If the engineer correctly understands that the input is stochastic, then she inverts each observed needle position to obtain the corresponding input value, thus eventually learning the true input distribution. But what if the engineer incorrectly anticipates a deterministic input and attributes the tremble of the needle to zero-mean measurement noise? Such an engineer must conclude that the deterministic input generates the average needle position. Her input estimate is the certainty equivalent of the input distribution under a Bernoulli utility function equal to the meter’s non-linear measurement function.

Our results on the behavioral implications of the perception strategy are analogous to the plight of the misspecified engineer. For simplicity, consider a DM who incorrectly anticipates a riskless lottery that pays the same reward in all states. Like the engineer who incorrectly anticipates a deterministic input, this DM estimates a single reward value, the perturbed encoding of which supposedly generated her perception data. The maximum-likelihood estimate of the encoded value of this single reward is the average of all the observed signals.

As the sample size diverges, the estimate converges to a convex combination of the encoded values of the states' true rewards, where the weight on each state is its sampling frequency. Hence, the DM's estimate of the lottery value converges to the certainty equivalent of the lottery evaluated with a Bernoulli utility function equal to the encoding function and subjective probabilities equal to the sampling frequencies.

We provide two extensions that bridge the gap between the extreme cases of a correctly specified DM who anticipates all possible risk and a misspecified DM who anticipates no risk at all. In our first approach, the DM is aware that she may face risk but uses a coarse partitional model of the true state space, much like Savage's (1954) decision-maker employing a small-world model of the grand world. The finest partition corresponds to the correctly specified DM, while the coarsest partition corresponds to the DM who anticipates no risk. There are various reasons why a DM might employ a coarse model. She might have evolved in a simple environment and the complexity of the environment might have increased, making previously payoff-irrelevant contingencies relevant, without the DM adapting to the change. For instance, the financial expert when analysing a new asset may initially estimate its expected performance in a coarse model that fails to include all economic contingencies that are relevant to that asset. Alternatively, the DM might have been framed to believe that the decision problem involves less risk than it does (by a car dealer, for example).

We find that, in the limit of nearly complete information, the coarse DM makes risk-neutral choices whenever she faces risk that is measurable with respect to her partition. But, whenever she faces a lottery that is not measurable with respect to her partition, she makes a biased choice even as her perception data become rich. She treats the lottery as if she had risk-attitudes implied by her perception strategy towards those elements of the risk that she does not comprehend, and is risk-neutral with respect to those elements of the risk that she does comprehend.

In our second approach, the DM anticipates some risk but finds large risks unlikely. We formalize this by taking a joint limit in which perception data become rich and the prior reward distribution gradually concentrates on the set of riskless lotteries. Perception distortions therefore remain large relative to the level of perceptual discrimination required in typical decision problems. We find risk attitudes akin to those of the DM who does not anticipate any risk at all. We then study comparative statics of these risk attitudes by varying the relative speed at which the two limits are taken. Within the parametrization we examine, choice becomes more risk-neutral when the DM anticipates larger risk a priori. In the context of Rabin's (2000) paradox, this implies that framing a decision problem as one which features high risk attenuates the DM's risk preferences. The DM also becomes more risk-neutral when she collects more data. Thus, the model predicts that risk attitudes are

induced under time pressure, mirroring the observation of Kahneman (2011) that prospect theory applies to fast instinctive decisions rather than to slow deliberative choices.

2 Related Literature

We build on a rich literature in neuroscience and economics. To our knowledge, our paper is the first to make two distinct contributions. First, we jointly optimize both encoding of the lottery rewards and their sampling frequencies. Second, we clarify the essential role of misspecification for behavioral consequences of any perception strategy when stakes are large relative to perception frictions.

Our work derives ultimately from psychophysics, a field that originated in Fechner’s (1860) study of stochastic perceptual comparisons based on Weber’s data.¹ A large literature in brain sciences and psychology views perception as information processing via a limited channel and studies the optimal encoding of stimuli for a given channel capacity. Laughlin (1981) derives and tests the hypothesis that optimal neural encoding under an information-theoretic objective encodes random stimuli with neural activities proportional to their cumulative distribution values. This implies S-shaped encoding for unimodal stimulus densities.²

Neuroscience studies encoding adaptations under various optimization objectives such as maximization of mutual information between the stimulus and its perception, maximization of Fisher information, or minimization of the mean squared error of perception.³ Economics can help here by providing microfoundations for the most appropriate optimization objective for perceptions related to choice. Robson (2001) has studied encoding of rewards that minimizes the probability of making a wrong choice and has shown that, in the limit of vanishing perception frictions, the optimal encoding function likewise coincides with the cumulative distribution function of rewards in the decision environment. Netzer (2009) has studied maximization of the expected chosen reward, an objective rooted in the instrumental approach of economics to information. The optimal encoding function still tracks the cumulative distribution function but is straightened. Schaffner et al. (2021) report that the optimal encoding function as in Netzer provides a better fit to neural data than do encodings derived under competing objectives.

These models study choices over riskless prizes and thus the derived encoding functions are not directly relevant to choices over gambles. Indeed, encoding functions are often inter-

¹Woodford (2020) provides a review of psychophysics from an economics perspective.

²See Attneave (1954) and Barlow (1961) for early contributions and Heng et al. (2020) for recent work.

³See e.g. Bethge et al. (2002) and Wang et al. (2016).

preted as hedonic anticipatory utilities rather than as Bernoulli utilities in that literature.⁴ We extend Netzer’s instrumental approach to choices over gambles, finding a connection to one of the above reduced-form objectives.⁵ That is, in the limit with rich perception data, maximization of the expected chosen reward is equivalent to minimization of the expected mean squared error in the perceived lottery value, where the expectation is over all decision problems with a tie. This conditioning on ties not only generates the better fit of the optimal encoding function documented by Schaffner et al. (2021), but is also crucial for the result of optimal oversampling of low-probability contingencies. Oversampling would not arise under reduced-form objectives that maximize unconditional measures of precision.⁶

Some recent papers study risk attitudes stemming from reward encoding in the presence of noise. Khaw et al. (2018) show theoretically and verify experimentally that exogenous logarithmic stochastic encoding and Bayesian decoding generates risk attitudes in an effect akin to reversion to the mean. Vieider (2021) proposes a model in which probabilities are also encoded in an exogenous logarithmic way and establishes a connection to stochastic prospect theory. Frydman and Jin (2019) and Juechems et al. (2021) allow for optimal encoding of the lottery reward and show both theoretically and experimentally that this encoding adapts to the distribution of the decision problems and that the adaptation affects choice. Relative to these papers, we analyze optimal encoding of rewards alongside optimal treatment of probabilities. We also differ in the proposed source of behavioral distortions. The discussed models assume well-specified learning, and thus they approximate the frictionless benchmark when noise becomes small. We focus on the limit of small encoding noise right away. This focus uncovers a novel connection between coding and behavior. While the impact of coding on behavior must necessarily vanish when the decoding model is well-specified, as in the previous literature, the implications for behavior remain substantial if the cognitive model used for decoding oversimplifies the structure of the risk. We connect perception to classical representations using a Bernoulli utility function and subjective probability weights.

We apply the statistical results of Berk (1966) and White (1982) on asymptotic outcomes

⁴See also Rayo and Becker (2007). Robson et al. (2021) is a dynamic version of Robson (2001) and Netzer (2009) that captures low-rationality, real-time adaptation of a hedonic utility function used to make ultimately deterministic choices. Friedman (1989) is an early approach dealing with gambles.

⁵Our model differs from Robson (2001) and Netzer (2009) concerning the perception friction. Those papers model frictions as minimal just noticeable differences, while here we rely on the modeling framework of Thurstone (1927) who hypothesized that perception is a Gaussian perturbation of an encoded stimulus. Payzan-LeNestour and Woodford (2021) have shown that the Gaussian approach yields the same limiting results as in Robson (2001) and Netzer (2009).

⁶Herold and Netzer (2015) derive probability weighting as the optimal correction for an exogenous distortive S-shaped value function, and Steiner and Stewart (2016) find probability weighting to be an optimal correction for naive noisy information processing. Lieder et al. (2017) argue that a contingency should be oversampled if it has extreme payoff consequences and decisions are based on a small sample. The present paper derives both S-shaped encoding and low-probability over-sampling in a joint optimization.

of misspecified Bayesian and maximum-likelihood estimation, respectively. The concept of Berk-Nash equilibrium in Esponda and Pouzo (2016) is defined as a fixed point of misspecified learning. This has motivated a renewed interest in misspecification across economics. Heidhues et al. (2018) characterize a vicious circle of overconfident learning, Molavi (2019) studies the macroeconomic consequences of misspecification, Frick et al. (2021) rank the short- and long-run costs of various forms of misspecification, and Eliaz and Spiegel (2020) focus on political-economy consequences of misspecification. We study the interplay of encoding and misspecified decoding of rewards. In his discussion of small-world models, Savage (1954) argues that a coarse representation of the complex grand world does not necessarily distort behavior. This is true whenever the subjective values assigned to the elements of a coarse state space partition are correct averages of the true rewards within each element. Our approach departs from Savage in that we explicitly model the process of learning about rewards. We argue that the DM is unlikely to learn the correct average rewards for each element of her partition. If she learns within the small-world model, then, instead of the average reward, her estimate converges to the certainty equivalent under her encoding function and subjective probabilities equal to her sampling frequencies.

Salant and Rubinstein (2008) and Bernheim and Rangel (2009) provide a revealed-preference theory of the behavioral and welfare implications of frames – payoff-irrelevant aspects of decision problems. We provide an account of how a specific frame – anticipation of the risk structure – affects choice and welfare. As in Kahneman, Wakker, and Sarin (1997), our model implies a distinction between decision and welfare utilities. In the case of the misspecified DM, the gap between the decision utility that she anticipates the lottery to pay and welfare utility – the true expected lottery reward – may be large. Our model facilitates an analysis of systematic mistakes in decision making as outlined in Koszegi and Rabin (2008) and, for the case of framing effects, Benkert and Netzer (2018).

3 Decision Process

There is a fixed set of states of the world $i \in \{1, \dots, I\}$, $I \geq 1$, where each state i has a fixed positive probability p_i . The DM chooses between a safe option of value s and a lottery that pays a reward $r_i \in [\underline{r}, \bar{r}]$ in each state i , where $\underline{r} < \bar{r}$ are arbitrary bounds. The lottery rewards and the safe option value are generated randomly.⁷ The DM observes the value of the safe option but faces frictions in the perception of the lottery rewards. We let $\mathbf{r} = (r_i)_i \in [\underline{r}, \bar{r}]^I$ denote the vector of rewards and simply refer to it as the *lottery*. The pair (\mathbf{r}, s) is the *decision problem*.

⁷Our results continue to hold when the value of the safe option is held fixed.

The goal of the DM is to choose the lottery if and only if its expected value $r = \sum_i p_i r_i$ exceeds s . This risk-neutrality with respect to rewards is an implicit assumption on the units of measurement in which the rewards are expressed. For instance, the rewards might be an appropriate concave function of monetary prizes if the DM chooses among monetary lotteries and money has diminishing returns.

The DM estimates the unknown lottery \mathbf{r} from a sequence of n signals, where each signal is a monotone transformation of one of the rewards perturbed with additive noise: she observes signals $x_k = (\hat{m}_k, i_k)$, $k = 1, \dots, n$. We refer to the first component, \hat{m}_k , as the *perturbed message*. The second component, i_k , indicates the state the message \hat{m}_k pertains to. Each perturbed message is generated by encoding the reward r_{i_k} in state i_k into *unperturbed message* $m(r_{i_k})$ and by perturbing it to $\hat{m}_k = m(r_{i_k}) + \hat{\varepsilon}_k$, where the noise term $\hat{\varepsilon}_k$ is independently and identically distributed (iid) standard normal.⁸ The sampled state i_k is one of the states $i = 1, \dots, I$, iid across k with positive probabilities π_i . The function $m : [\underline{r}, \bar{r}] \rightarrow [\underline{m}, \bar{m}]$ is strictly increasing and continuously differentiable; we refer to it as the *encoding function*. We dub π_i as *sampling frequencies* and refer to $(m(\cdot), (\pi_i)_i)$ as the *perception strategy*. The size of the sample, n , is exogenous.

After she has observed the n signals, the DM forms an estimate q_n of the lottery's value and chooses the lottery if and only if $q_n > s$. We consider both maximum-likelihood and Bayesian estimators, $q_n = q_n^{ML}$ or $q_n = q_n^B$. In the first case, the DM is endowed with a compact set $\mathcal{A} \subseteq [\underline{r}, \bar{r}]^I$ of lotteries she anticipates and concludes that she has encountered the lottery

$$\mathbf{q}_n^{ML} \in \arg \max_{\mathbf{r}' \in \mathcal{A}} \prod_{k=1}^n \varphi(\hat{m}_k - m(r'_{i_k}))$$

that maximizes the likelihood of the observed signals, where φ is the standard normal density. Finally, she sets $q_n^{ML} = \sum_i p_i q_{in}^{ML}$.⁹ In the second case, the DM is endowed with a prior belief over \mathcal{A} and sets $q_n^B = \mathbb{E}[\sum_i p_i r_i \mid (x_k)_{k=1}^n]$ as the posterior expected lottery value. Both these specifications will lead to same conclusions as n diverges since the impact of the DM's prior becomes negligible in this limit.

We study decision-makers who may employ simplifying models of risk in the spirit of the *small world* of Savage (1954). The DM anticipates, rightly or wrongly, distinctions among some of the states of the world to be payoff-irrelevant. Let \mathcal{P} be a partition of the set of all the states $\{1, \dots, I\}$. The DM anticipates that $r_i = r_j$ for all pairs of states $i, j \in J$ that

⁸Our results in Section 4 on the limit loss and the optimal perception strategy extend to general noise distributions. We use the Gaussian assumption in Section 5, where it yields a tractable form of the Kullback-Leibler divergence and intuitive outcomes of misspecified learning.

⁹The maximum-likelihood estimate exists since \mathcal{A} is compact. It is unique for the specifications below.

belong to a same element J of the partition \mathcal{P} . That is, she anticipates lotteries from a set

$$\mathcal{A}_{\mathcal{P}} = \left\{ \mathbf{r} \in [\underline{r}, \bar{r}]^I : r_i = r_{i'} \text{ for all } i, i', J \text{ such that } i, i' \in J, J \in \mathcal{P} \right\}. \quad (1)$$

For instance, if $\mathcal{P} = \{\{1, \dots, I\}\}$ is the coarsest partition, then the DM anticipates only degenerate lotteries that pay a same reward in all states. We refer to such lotteries as *riskless* and call other lotteries *risky*. If, on the other extreme, $\mathcal{P} = \{\{1\}, \dots, \{I\}\}$ is the finest partition, then the DM anticipates that any reward vector is possible and $\mathcal{A}_{\mathcal{P}} = [\underline{r}, \bar{r}]^I$.

4 Optimal Perception

The perception strategy needs to adapt to the prevailing statistical circumstances if it is to allocate attention efficiently. An increase of the sampling frequency of a state increases the DM's attention to its reward, but reduces attention to the rewards in other states. Similarly, making the encoding function steep in a neighborhood of a reward value reduces noise in this neighborhood but entails increased noise elsewhere.

We denote the state space partition that the DM employs during the adaptation stage by \mathcal{J} . That is, the DM anticipates lotteries from $\mathcal{A}_{\mathcal{J}}$ where each element of partition \mathcal{J} specifies a set of states that the DM deems as payoff-equivalent. Since the distinction between states within each $J \in \mathcal{J}$ is redundant, we treat J as an index of a state, refer to the rewards in states $i \in J$ simply as r_J , and model the whole lottery $\mathbf{r} = (r_J)_{J \in \mathcal{J}}$ as having $|\mathcal{J}|$ rewards, each with probability $p_J = \sum_{i \in J} p_i$. A perception strategy consists of the increasing encoding function $m(\cdot)$ and interior sampling frequencies $(\pi_J)_J \in \Delta(\mathcal{J})$.¹⁰ In Section 5, we will use this notation to study a DM whose model is a misspecified small-world model of the grand world, for instance because the world became more complex after adaptation but before choice.

The DM optimizes her perception strategy ex ante for a given distribution of decision problems. Specifically, the rewards r_J are iid with a continuous density h , and the safe option s is drawn from a continuous density h_s independently of the lottery rewards; both densities have supports $[\underline{r}, \bar{r}]$.¹¹ We characterize the expected loss for general perception strategies for diverging n in the next subsection and then solve for the loss minimizing strategy in Subsection 4.2.

¹⁰Assuming interior sampling frequencies (when $|\mathcal{J}| > 1$) is without loss, because in the limit when the number of signals grows it is optimal to gather at least some information about the rewards in each state.

¹¹Since s can have a distinct density from that of r_J , the safe option may, for example, capture in reduced form the choice of an alternative lottery with each of its rewards drawn from h .

4.1 Objective

We take the number n of signals to be large and abstract from uncertainty over the number of perturbed messages sampled for each state and from divisibility issues. That is, we suppose the number of messages sampled for each state $J \in \mathcal{J}$ is precisely $\pi_J n$. We let $m_{J,n}$ be the average of the perturbed messages sampled for state J . Then, $m_{J,n} - m(r_J)$ is normally distributed with mean 0 and variance $1/(n\pi_J)$, for each given value of r_J . Since the signal errors are Gaussian, the vector of average perturbed messages, $\mathbf{m}_n = (m_{J,n})_J$, is a sufficient statistic for the lottery rewards.

For $z \in \{B, ML\}$, let q_n^z be the Bayesian and maximum-likelihood estimator of the lottery value and let

$$L^z(n) = \mathbb{E} \left[\max\{r, s\} - \mathbb{1}_{q_n^z > s} r - \mathbb{1}_{q_n^z \leq s} s \right]$$

be its ex ante expected loss relative to choice under complete information; the expectation is over \mathbf{r} , s and q_n^z .

Proposition 1. *Assume the encoding function m is continuously differentiable, the reward density h is continuous, and the density of the safe option h_s is continuously differentiable. Then, the Bayesian and maximum-likelihood estimators generate the same asymptotic loss*

$$\lim_{n \rightarrow \infty} nL^z(n) = \frac{1}{2} \mathbb{E} \left[h_s(r) \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J)} \right] \text{ for } z \in \{B, ML\}. \quad (2)$$

See Appendix A for the proof. The difference between the Bayesian and maximum-likelihood estimators asymptotically vanishes because the prior information has negligible impact on the Bayesian DM who receives many signals. The limit loss characterization in (2) has an intuitive interpretation. It is the mean squared error (MSE) in the perception of the lottery value (multiplied by $n/2$) integrated over all decision problems in which the true lottery value r ties with s . The conditioning on the tie arises because the likelihood of large perception errors vanishes with increasing n , and small perception errors distort choice only in decision problems in which an approximate tie arises. In the limit, the set of decision problems in which perception errors have nontrivial behavioral consequences approaches the set of problems with exact ties.

To understand the relevance of the MSE for loss, fix the true and perceived lottery values to be r and q_n^z , respectively. The perception error distorts choice and causes loss if and only if the safe option s attains a value between r and q_n^z . When n is large, and hence the error is small, this occurs with approximate probability $h_s(r)|q_n^z - r|$. Conditional on the choice being distorted like this, the expected loss is approximately $|q_n^z - r|/2$ since s is approximately

uniformly distributed between r and q_n^z . Hence the overall loss over all s is approximately $h_s(r) (q_n^z - r)^2 / 2$. Taking the expectation of $(q_n^z - r)^2$ with respect to q_n^z yields the MSE of the value estimate for a given lottery, and taking the expectation with respect to the lottery gives an average over all decision problems with a tie, $s = r$.

To understand the expression in (2) in detail, consider the maximum-likelihood estimator; the Bayesian estimator differs only by a negligible term. The maximum-likelihood estimate of the reward r_J is

$$q_{J,n}^{ML} = m^{-1}(m_{J,n}), \quad (3)$$

and, since $m_{J,n} \sim \mathcal{N}(m(r_J), 1/(n\pi_J))$, the MSE of $q_{J,n}^{ML}$ is approximately $1/(n\pi_J m'^2(r_J))$.¹² The MSE of the value estimate is then approximately

$$\frac{1}{n} \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J)}.$$

The MSE and hence the loss go to zero as $n \rightarrow \infty$, for any perception strategy. However, the perception strategy influences how fast the loss vanishes. Motivated by the characterization from (2), we define the *information-processing problem* as follows.

$$\min_{m'(\cdot) > 0, (\pi_J)_J > 0} \quad \mathbb{E} \left[\sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J)} \mid r = s \right] \quad (4)$$

$$\text{s.t.:} \quad \int_{\underline{r}}^{\bar{r}} m'(\tilde{r}) d\tilde{r} \leq \bar{m} - \underline{m} \quad (5)$$

$$\sum_{J \in \mathcal{J}} \pi_J = 1. \quad (6)$$

The objective in (4) equals the asymptotic loss characterized in (2), up to a factor that is independent of the perception strategy.¹³ We let the DM control the derivative $m'(\cdot)$ and restrict it to be positive – this restricts the encoding function to be increasing and differentiable. Constraint (5) is implied by the finite range of the encoding function – the encoding function cannot be steep everywhere. Constraint (6) together with the restriction to positive sampling frequencies requires $(\pi_J)_J$ to be a probability distribution over \mathcal{J} – the

¹²Equation (3) holds if $m_{J,n} \in [\underline{m}, \bar{m}]$. If $m_{J,n} < \underline{m}$ or $m_{J,n} > \bar{m}$, then the maximum-likelihood estimate of r_J is \underline{r} or \bar{r} , respectively. Note, that $P(m_{J,n} \in [\underline{m}, \bar{m}]) \rightarrow 1$ as n diverges for all $r_J \in (\underline{r}, \bar{r})$.

¹³This factor is two divided by the ex ante likelihood of a tie. A special case in which conditioning on ties can be ignored is when s is uniformly distributed on $[\underline{r}, \bar{r}]$, because conditional MSE and unconditional MSE are then identical up to a constant factor. In that case, our information-processing problem implies the minimization of the unconditional MSE, an objective assumed for example by Woodford (2012).

DM must also treat sampling frequencies as a scarce resource.

4.2 Optimization

We say that the perception strategy $(m(\cdot), (\pi_J)_J)$ is optimal if $(m'(\cdot), (\pi_J)_J)$ solves the information-processing problem. To describe the optimal strategy, we denote by

$$h_J(\tilde{r}) = h(\tilde{r}) \frac{\mathbb{E}[h_s(r) | r_J = \tilde{r}]}{\mathbb{E}[h_s(r)]}$$

the density of reward r_J in state J conditional on a tie between the lottery value and the safe option (the expectations are over \mathbf{r}). Observe that each h_J is continuous if both h and h_s are continuous,¹⁴ which we assume in the following.

Proposition 2. *There is a unique optimal perception strategy. It has the properties:*

1. *The optimal encoding function satisfies, for all $\tilde{r} \in [\underline{r}, \bar{r}]$,*

$$m'(\tilde{r}) \propto \left(\sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J} h_J(\tilde{r}) \right)^{\frac{1}{3}}. \quad (7)$$

2. *The optimal sampling frequencies satisfy, for all $J, J' \in \mathcal{J}$,*

$$\left(\frac{p_J}{\pi_J} \right)^2 \mathbb{E} \left[\frac{1}{m'^2(r_J)} \mid r = s \right] = \left(\frac{p_{J'}}{\pi_{J'}} \right)^2 \mathbb{E} \left[\frac{1}{m'^2(r_{J'})} \mid r = s \right]. \quad (8)$$

The proof of Proposition 2 in Appendix B follows from first-order conditions. The first-order condition for the slope $m'(\tilde{r})$ of the encoding function is

$$2 \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^3(\tilde{r})} h_J(\tilde{r}) = \lambda \quad (9)$$

for each reward value \tilde{r} , where λ is the shadow price of constraint (5). The left-hand side of (9) is the marginal benefit of an increase in the slope $m'(\tilde{r})$ at the reward value \tilde{r} . Such an increase reduces the DM's MSE in her perception of the lottery value if the reward r_J attains the value \tilde{r} for one of the states $J \in \mathcal{J}$. This marginal reduction affects her choice if the value of the lottery r ties with s . Each summand on the left-hand side is proportional to the marginal reduction of the MSE multiplied by the likelihood that $r_J = \tilde{r}$ and that

¹⁴Since h_s is continuous on a compact interval, it is uniformly continuous, and thus the function $\tilde{r} \mapsto \mathbb{E}[h_s(r) | r_J = \tilde{r}]$ is continuous. Thus, h_J is continuous.

$r = s$. The constraint (5) implies that, at the optimum, the marginal benefit of a slope increase is equal across all reward values \tilde{r} . Expressing $m'(\tilde{r})$ from (9) gives the explicit solution (7) in the proposition. This solution generalizes Netzer (2009). When $|\mathcal{J}| = 1$, then our DM chooses between two riskless rewards r and s . Both Netzer and we find that when r and s are independently drawn from a same density h , then the slope of the optimal encoding function is proportional to $h^{2/3}(r)$. To see this in our framework, note that the reward density conditional on a tie is proportional to $h^2(r)$ for $|\mathcal{J}| = 1$ and the result then follows from (7).

Property (8) reflects that, at the optimum, the marginal benefit of increasing the sampling frequency must be equal across all states J , because of constraint (6). This is, broadly speaking, achieved by matching the probability p_J of each state with its sampling frequency π_J , because more precise information is more valuable for states which arise with higher probability and hence are more relevant for the lottery value. Relative to the true probabilities, however, the DM wishes to over-sample states about whose rewards she expects to be poorly informed. Recall that the DM measures reward r_J relatively poorly if the slope $m'(r_J)$ is low. According to (8), the DM therefore chooses her (squared) sampling frequencies to match (squared) modified probabilities, where the modifying factor is the expectation of $1/m^2(r_J)$ and reflects how poorly the DM expects to be informed about the reward in state J . The expectation is again conditional on a tie because a marginal change of the sampling frequency affects choice only at ties.

A density $f(x)$ on $[\underline{r}, \bar{r}]$ is *unimodal* and *symmetric* around the mode $r_m = (\underline{r} + \bar{r})/2$ if it is strictly decreasing on $(r_m, \bar{r}]$ and $f(r_m + x) = f(r_m - x)$ for all x . Symmetry implies that unimodality is preserved by summation. We obtain the following proposition for the case of unimodal and symmetric reward densities.¹⁵

Proposition 3. *If the densities h and h_s are unimodal and symmetric, then the optimal perception strategy has the properties:*

1. *The optimal encoding function is S-shaped. It is convex below and concave above r_m .*
2. *The DM over-samples low-probability states. For any two states $J, J' \in \mathcal{J}$ such that $p_J < p_{J'}$, it holds that $\frac{\pi_J}{p_J} > \frac{\pi_{J'}}{p_{J'}}$. In particular, when there are two states, then $\pi_J > p_J$ for the state with probability $p_J < 1/2$ and vice versa for the high-probability state.*

We show in Appendix B.2 that the density of the reward conditional on a tie is, for each state, unimodal with the same mode as the unconditional reward density. The solution

¹⁵If the safe option is the value of an alternative lottery with rewards drawn from h , as discussed in footnote 11, then unimodality and symmetry of h naturally implies unimodality and symmetry of h_s .

(7) thus implies that the optimal slope is proportional to a monotone transformation of a sum of unimodal functions that all have their maxima at the unconditional reward mode, establishing the first claim of the proposition.

While rewards are iid across the states unconditionally, conditional on a tie they are no longer identically distributed. The tie condition $\sum_J p_J r_J = s$ is relatively uninformative about rewards in low-probability states, and hence the conditional reward distributions for the low-probability states are more spread-out compared to the high-probability states (see Appendix B.2). Because the optimal m is relatively flat at tail rewards, if the DM sampled in proportion with the states' probabilities then she would in expectation end up poorly informed about the spread-out rewards of low-probability states, and hence the marginal benefit of an additional signal would be larger for those than for the other states. She optimally compensates by over-sampling the low-probability states relative to proportional sampling.¹⁶ Optimal over-sampling thus arises from our microfoundation of objective (4). Had the DM minimized the unconditional MSE, the effect would not arise. By taking the instrumental perspective that focuses on the payoff consequences of perception errors in choice problems, we obtain an objective that conditions on ties and induces over-sampling as the optimal adaptation.

4.3 Extension

So far we have treated the partition \mathcal{J} as fixed and known at the stage of optimization. Hence the optimal perception strategy depends on the partition. Furthermore, optimal sampling frequencies are only determined for the elements of the partition but not separately for each of the states within an element. In Appendix B.3, we provide an extension of the model in which optimization of the encoding function m and sampling frequencies π_i for all states $i = 1, \dots, I$ takes place before the partition is realized. This is relevant when evolutionary forces select the perception strategy at a slow pace and can condition only on coarse information about the partition, while the DM obtains more information before sampling takes place. We generalize the optimality conditions from Proposition 2 and show that, under the assumptions of Proposition 3, an S-shaped encoding function is still optimal. The qualitative intuition for the optimal sampling frequencies is also still valid: it is optimal to over-sample states of the world that are typically included in low-probability events.

¹⁶This argument relies on the rewards for all states being encoded with the same encoding function. The property of S-shaped encoding would still hold with state-dependent encoding functions, but the optimal sampling frequencies would be more difficult to derive. They would, for example, depend on whether the range constraint (5) applies to each state separately or across states, i.e., on whether slope can be transferred across states or not.

5 Behavior

The implications of the perception strategy for behavior depend on the DM’s degree of understanding of the risk. Consider the example from Savage (1954) mentioned earlier. The DM is contemplating the purchase of a convertible car for price s . The payoff from the purchase depends on the random weather; it is r_1 if the car is driven in rainy conditions and r_2 for sunny conditions. The upcoming weather is unknown, making the purchase a binary lottery. Let the probabilities of either weather type be one half.

The DM learns the values of r_1 and r_2 by sampling n signals. For each $k = 1, \dots, n$, she observes the weather $i_k \in \{1, 2\}$ and a message $\hat{m}_k = m(r_{i_k}) + \hat{\varepsilon}_k$ where m is the encoding function and the $\hat{\varepsilon}_k$ are iid standard normal. The sampling frequency of each weather condition is for now assumed to be one half, thus matching the actual probabilities. Each signal might derive from the DM’s own experience with a convertible, the experience of her peers, information provided by the car dealer, etc.

Consider two varieties of DM – *fine* and *coarse* – who differ in their anticipation of the risk structure. The fine DM knows that the weather is payoff-relevant and hence anticipates that the purchase will lead to one of two possibly distinct reward values (r_1, r_2) . The coarse DM employs a small-world model: she anticipates, as in Savage’s example, that the convertible will lead to “definite and sure enjoyments”, so she anticipates a degenerate lottery (r, r) .

Their distinct models of risk lead the two DMs to distinct conclusions even when they employ the same perception strategy and observe identical data. The fine DM asymptotically learns $m(r_i)$ for $i = 1, 2$ from the empirical distribution of the perturbed messages, inverts the encoding function and learns the true reward pair. Her estimate of the expected reward thus converges to the true expected reward and she makes the risk-neutral choice. See the left-hand graph in Figure 1.

The coarse DM observes the same empirical signal distribution but, since she omits the weather from her model of risk, she seeks a single message which best accounts for all the observed signals. For Gaussian additive errors, the single message that maximizes the likelihood of the observed data is the empirical average message, which almost surely converges to $(m(r_1) + m(r_2))/2$. Hence, the DM’s asymptotic estimate of the reward from driving the convertible is the certainty equivalent of the risky reward under the Bernoulli utility $u(\cdot) = m(\cdot)$ and equal probabilities. See the right-hand graph in Figure 1.

There are various paths that could have led the fine and the coarse DMs to their respective decision procedures. They could have evolved in a simple environment in which all the lotteries were measurable with respect to the coarsest partition $\mathcal{J} = \{\{1, 2\}\}$ of the set of states. Afterwards, their environments became more complex so they currently encounter




Figure 1: Asymptotic estimated lottery value q of the fine (left) and the coarse (right) DMs.

lotteries with $r_1 \neq r_2$. The fine DM has refined her anticipation and understands that she may now face a risky lottery. In contrast, the coarse DM has not made such an adjustment and continues to anticipate riskless lotteries only. It is plausible that real-world decision-makers are sometimes not aware of all contingencies that affect their payoffs. Below we will provide an example of a financial investor who omits a relevant variable from her econometric model of the return of an asset. Alternatively, both DMs may have evolved in a risky environment with partition $\mathcal{J} = \{\{1\}, \{2\}\}$ or in which this partition was at least possible with positive probability. Afterwards, the coarse DM was (incorrectly) assured that her next lottery will be riskless (possibly by a strategically interested party), while the fine DM was not told this. Finally, both DM's may know that they encounter a risky lottery but the coarse DM has chosen the coarse estimation procedure due to its simplicity. The coarse procedure consists of applying the inverse encoding function to the average of all perturbed messages, whereas the procedure of the fine DM requires applying the coarse procedure to each state of the world separately and then computing the lottery value.

This section takes the DM's perception strategy as given; it could have been optimized as in Section 4 or established by any different process. Subsection 5.1 extends the present binary example to arbitrary lotteries and sampling frequencies. A further generalization in Subsection 5.2 considers a DM who employs an arbitrary partitional model of risk; such a DM has some but only partial awareness of the risk she faces. Subsection 5.3 then focuses on a DM who anticipates risk but believes that large differences between rewards across the states of the world are a priori unlikely. As in the case of the DM who anticipates no risk, this generates non-trivial risk attitudes.

5.1 Surprising Risk

We characterize here the behavior of a DM who has not anticipated any risk. She anticipates a lottery from the set

$$\mathcal{A} = \{\mathbf{r} \in [\underline{r}, \bar{r}]^I : r_i = r_j \text{ for all states } i, j\}.$$

After she encounters a lottery, she observes data generated by her perception strategy, forms the maximum-likelihood or Bayesian estimate of the encountered lottery from \mathcal{A} , and chooses the lottery if and only if its estimated value exceeds s . The DM learns in a misspecified model – she may encounter an unanticipated risky lottery.

To describe her behavior, we say that the DM’s choice is represented by a Bernoulli utility $u(\cdot)$ and probabilities $(\rho_i)_{i=1}^I$ if in each decision problem (\mathbf{r}, s) such that

$$\sum_{i=1}^I \rho_i u(r_i) > [<] u(s),$$

the probability that the DM chooses the lottery \mathbf{r} converges to 1 [0] as $n \rightarrow \infty$.¹⁷

Proposition 4. *Let the DM form the maximum-likelihood or Bayesian estimate of the lottery. When she anticipates a riskless lottery, the DM’s choice is represented by a Bernoulli utility equal to the encoding function, $u(\cdot) = m(\cdot)$, and probabilities given by the sampling frequencies, $\rho_i = \pi_i$ for $i = 1, \dots, I$.*

The proposition is a special case of Proposition 5 which we prove in Appendix C. It follows from the result on misspecified maximum-likelihood estimation by White (1982) and from Berk (1966) for the Bayesian DM. These authors let an agent observe n iid signals from a signal density and form the estimate from a set of hypothesized signal densities that may fail to include the true density. They prove that the estimate almost surely converges to the minimizer of the Kullback-Leibler divergence from the true signal density as n diverges (if the minimizer is unique).

To apply White’s and Berk’s results in our setting, consider a DM who encounters a lottery \mathbf{r} . She observes the empirical distribution of approximately $\pi_i n$ signals drawn iid from $\mathcal{N}(m(r_i), 1)$ for each state i . Since the DM has anticipated a riskless lottery, she forms an estimate of a single unperturbed message m_n , a perturbation of which has generated the observed data. White’s and Berk’s results imply that $m^* = \lim_{n \rightarrow \infty} m_n$ almost surely minimizes the Kullback-Leibler divergence from the true signal density. For Gaussian errors,

¹⁷The probability is evaluated with respect to the stochastic signal sequence $(\hat{m}_k, i_k)_{k=1}^n$.

this implies $m^* = \sum_i \pi_i m(r_i)$ almost surely. Thus, the DM's estimate of the lottery value almost surely converges to the "certainty equivalent" $m^{-1} \left(\sum_{i=1}^I \pi_i m(r_i) \right)$.

The behavior of the DM who anticipates a riskless lottery is governed by the sampling frequencies rather than by the true probabilities. Indeed, this DM believes that the true probabilities are payoff-irrelevant. In contrast, the sampling frequencies govern the proportions of her data generated for each state and hence her estimate of the encoded riskless reward she thinks she has encountered.

Example (estimating financial returns): An investor chooses between a safe asset with return s and a risky asset with return $\rho(\mathbf{x}, \mathbf{y})$ that depends on vectors of variables \mathbf{x} and \mathbf{y} . She employs a misspecified model: she neglects the role of variables \mathbf{y} , believing that the return is $\tilde{\rho}(\mathbf{x})$ where $\tilde{\rho}(\cdot)$ is a simplified function she estimates. For example, she knows that the profit of a firm depends on prices and interest rate (\mathbf{x}) but is not aware of the firms' entire trade exposure and neglects the role of some exchange rates (\mathbf{y}). Given \mathbf{x} , let \mathbf{y} have conditional probability $g(\mathbf{y} | \mathbf{x})$. Thus, for each fixed value of \mathbf{x} the financial asset is a lottery where each state represents a particular value of \mathbf{y} and is assigned a return $\rho(\mathbf{x}, \mathbf{y})$ and a probability $g(\mathbf{y} | \mathbf{x})$. However, the investor attributes the variation of the return for fixed \mathbf{x} to noise and estimates $\tilde{\rho}(\mathbf{x})$ from signals $m(\rho(\mathbf{x}, \mathbf{y}_k)) + \hat{\varepsilon}_k$, $k = 1, \dots, n$. The conditional probability $\tilde{g}(\mathbf{y} | \mathbf{x})$ of observing a signal about the return for $\mathbf{y}_k = \mathbf{y}$ depends on the investor's sampling; if her sampling is representative, then $\tilde{g} = g$. By Proposition 4, when the number of signals diverges, the investor treats the asset for each \mathbf{x} as if she were an expected-utility maximizer with Bernoulli utility $u(\cdot) = m(\cdot)$ and probability $\tilde{g}(\mathbf{y} | \mathbf{x})$ assigned to each value of y .

5.2 Coarse Decision-Maker

Next, we study a DM who considers distinctions among some but not all states of the world payoff-relevant. She anticipates that all states in each element of a partition \mathcal{K} of the set of states pay the same reward. That is, she anticipates encountering a lottery from the set $\mathcal{A}_{\mathcal{K}}$ of lotteries measurable with respect to \mathcal{K} .

We say that the DM's choice has a *mixed representation* with Bernoulli utility $u(\cdot)$, probabilities $(\rho_i)_{i=1}^I$ and partition \mathcal{K} if the probability that she chooses lottery \mathbf{r} over the safe option s converges to 1 [0] in each decision problem (\mathbf{r}, s) such that

$$\sum_{J \in \mathcal{K}} \rho_J r_J^* > [<] s,$$

where $\rho_J = \sum_{i \in J} \rho_i$ and r_J^* is the certainty equivalent defined by

$$u(r_J^*) = \sum_{i \in J} \frac{\rho_i}{\rho_J} u(r_i)$$

for each $J \in \mathcal{K}$.

Let $J(i)$ be the element of partition \mathcal{K} that contains state i . Let $p_J = \sum_{i \in J} p_i$ be the overall probability of the states $i \in J$, and let $\pi_J = \sum_{i \in J} \pi_i$ be the overall sampling frequency for J .

Proposition 5. *The choice of the coarse DM who forms a maximum-likelihood or Bayesian estimate has a mixed representation with Bernoulli utility $u(\cdot) = m(\cdot)$ and probabilities $\rho_i = p_{J(i)} \frac{\pi_i}{\pi_{J(i)}}$ for $i = 1, \dots, I$.*

See Appendix C for the proof. In the limit, the DM chooses as if she was treating the lottery \mathbf{r} as a compound lottery in which each element J of the partition \mathcal{K} constitutes a sub-lottery and these sub-lotteries have probabilities p_J . She behaves as if she first reduced each sub-lottery to its certainty equivalent under the Bernoulli utility $u(\cdot) = m(\cdot)$ and subjective probabilities equal to the normalized sampling frequencies. After the reduction, she evaluates the overall lottery in a risk-neutral manner using the true probabilities of each J .

Example (estimating financial returns continued): Unlike in the previous version of this example, the investor does not observe \mathbf{x} (or \mathbf{y}) at the moment of choice. Instead, she observes a signal \mathbf{z} . Conditional on the observed value of \mathbf{z} , the asset is a lottery in which each state represents a realization of (\mathbf{x}, \mathbf{y}) with associated return $\rho(\mathbf{x}, \mathbf{y})$ and probability $g(\mathbf{x}, \mathbf{y} \mid \mathbf{z})$. Since the investor is unaware of \mathbf{y} 's influence on the return, she forms a coarse counterpart of this lottery in which each state represents a value of \mathbf{x} , paying $\tilde{\rho}(\mathbf{x})$ with probability $g(\mathbf{x} \mid \mathbf{z}) = \sum_{\mathbf{y}} g(\mathbf{x}, \mathbf{y} \mid \mathbf{z})$. For each value of \mathbf{x} , the investor forms the estimate of the return $\tilde{\rho}(\mathbf{x})$ given the data points $m(\rho(\mathbf{x}, \mathbf{y}_k)) + \hat{\varepsilon}_k$, where \mathbf{y}_k is drawn from $\tilde{g}(\mathbf{y}_k \mid \mathbf{x}, \mathbf{z})$. Again, $\tilde{g}(\mathbf{y} \mid \mathbf{x}, \mathbf{z})$ captures sampling. If sampling is untargeted, then $\tilde{g} = g$. After she forms the maximum-likelihood estimate $\hat{\rho}_n(\mathbf{x})$ for each value \mathbf{x} , she assigns the expected value $E[\hat{\rho}_n(\mathbf{x}) \mid \mathbf{z}]$ to the asset, where the expectation is with respect to the conditional density $g(\mathbf{x} \mid \mathbf{z})$. By Proposition 5, for each \mathbf{z} this investor values the asset as if she computed the certainty equivalent over $\rho(\mathbf{x}, \mathbf{y}) \mid (\mathbf{x}, \mathbf{z})$ for each (\mathbf{x}, \mathbf{z}) under Bernoulli utility $m(\cdot)$ and subjective probabilities $\tilde{g}(\mathbf{y} \mid \mathbf{x}, \mathbf{z})$, and then computed the risk-neutral value of the reduced lottery under the objective probabilities $g(\mathbf{x} \mid \mathbf{z})$. That is, the investor is risk-neutral with respect to the risk induced by stochastic $\mathbf{x} \mid \mathbf{z}$ that she comprehends but behaves as if she had non-trivial risk-attitudes with respect to the risk induced by stochastic $\mathbf{y} \mid (\mathbf{x}, \mathbf{z})$ that she does not comprehend.

If the DM actually encounters a lottery $\mathbf{r} \in \mathcal{A}_{\mathcal{K}}$ that she has anticipated, she learns in a correctly specified model. The asymptotic results for correctly specified learning of Wald (1949) for maximum-likelihood estimation and of Le Cam (1953) for Bayesian estimation imply that she correctly learns the encountered lottery as the number of signals diverges. In this case, our result implies that her perception strategy is irrelevant for her limit choice and she chooses in a risk neutral way.

Our predictions of the DM’s risk attitudes more generally depend on the combination of the adaptation experienced, as in Section 4, and her misapprehension of the lottery at the moment of choice. Recall that \mathcal{J} denotes the partition that the DM has employed during adaptation and partition \mathcal{K} specifies the DM’s anticipation of lotteries at the moment of choice; \mathcal{J} and \mathcal{K} may differ. The optimal encoding function is S-shaped regardless of the adaptation partition \mathcal{J} . Hence, we predict risk aversion (loving) for upper (lower) tail rewards with respect to the unanticipated risk under \mathcal{K} . We also predict relative sampling frequencies for states within the same element of partition \mathcal{K} whenever the distinction between these states was recognized during adaptation. If the DM was distinguishing these states under adaptation partition \mathcal{J} , or if adaptation occurred under sufficient uncertainty about the relevant partition (as in our extension in Subsection 4.3), we predict overweighting of low-probability events. For example, a DM who has adapted to a world in which weather condition is sometimes payoff-relevant, but was framed by a car dealer to ignore the role of the weather in the specific decision to purchase a convertible, may overweight the rather small probability of weather conditions suitable for driving open.¹⁸

5.3 Somewhat Surprising Risk

As a last extension of our model, we analyze a DM who deems risk a priori possible but unlikely. Her perception frictions are comparable in size to the risk that she typically expects to encounter. To this end, we study a limit in which the prior shrinks to the set of riskless lotteries as the amount of perception data diverges. We find perception distortions that are qualitatively similar to those from Subsection 5.1. Additionally, the approach makes predictions about the impact of framing and time pressure on risk-taking. Risk attitudes are attenuated by the anticipation of high risk or by rich perception data.

¹⁸If the DM has ignored distinction between two states already at the stage of adaptation, our results of Section 4 do not predict relative sampling frequencies. If sampling is representative, then the sampling frequencies coincide with their states’ objective probabilities. Any targeted sampling, for instance over-sampling salient contingencies, results in choice that assigns disproportional subjective probabilities to the over-sampled states. Starmer and Sugden (1993) report that a payoff-irrelevant split of an event increases the weight that lab subjects assign to this event. This effect arises for our coarse DM if splitting a contingency leads to it having a larger overall sampling frequency.

The DM of this subsection is Bayesian. Her prior density indexed by n is

$$\varrho_n(\mathbf{r}) = \varrho_n^0 \exp\left(-\frac{n}{2\Delta}\sigma^2(\mathbf{r})\right) \quad (10)$$

with support $[\underline{r}, \bar{r}]^I$, where $\sigma^2(\mathbf{r}) = \sum_{i=1}^I p_i(r_i - r)^2$ is the variance of the states' rewards and $r = \sum_i p_i r_i$ is the true lottery value as usual; ϱ_n^0 is the normalization factor. This prior is mostly concentrated on low-risk lotteries. For any fixed n , $\Delta > 0$ parameterizes the level of a priori anticipated risk. The index n has two roles. As n increases, risky lotteries become a priori less likely, approximating then the anticipation of the DM from Subsection 5.1. In addition to risk becoming less likely, the DM observes more data as n increases. She observes, for each state i , a sequence of $a\pi_i n$ messages equal to $m(r_i)$ perturbed with iid additive standard normal noise, where $(\pi_i)_{i=1}^I$ continues to denote the sampling frequencies. The parameter $a > 0$ captures attention span; the larger a is, the more signals the DM observes for each fixed n . The DM chooses the lottery \mathbf{r} over the safe option s if and only if the Bayesian posterior expected lottery value exceeds s .

To formulate the next result, we define a function $\mathbf{q}^* : [\underline{r}, \bar{r}]^I \rightarrow [\underline{r}, \bar{r}]^I$ as follows:

$$\mathbf{q}^*(\mathbf{r}) = \arg \min_{\mathbf{r}' \in [\underline{r}, \bar{r}]^I} \left\{ \frac{\sigma^2(\mathbf{r}')}{a\Delta} + \sum_{i=1}^I \pi_i (m(r'_i) - m(r_i))^2 \right\}. \quad (11)$$

We impose the regularity condition that the minimizer is unique. We refer to the posterior expectation $\mathbb{E}[\mathbf{r} \mid \mathbf{m}_n] \in [\underline{r}, \bar{r}]^I$ that the DM forms given the vector of the average perturbed messages \mathbf{m}_n as the Bayesian estimate of the lottery.

Proposition 6. *Suppose the DM has encountered lottery \mathbf{r} . The Bayesian estimate of the lottery converges to $\mathbf{q}^*(\mathbf{r})$ in probability as $n \rightarrow \infty$.*

See Appendix D for the proofs for this subsection. The asymptotic estimate $\mathbf{q}^*(\mathbf{r})$ of the lottery \mathbf{r} is a compromise lottery that is not too risky and does not generate messages too far from the true messages. When $a\Delta$ is small, then the DM anticipates relatively little risk and/or collects little perception data. Her best explanation of her perception data is a lottery that involves little risk. In the limit as $a\Delta \rightarrow 0$, the solution to (11) minimizes Kullback-Leibler divergence from the true lottery among the riskless lotteries, as in Subsection 5.1. When $a\Delta$ is large, then the DM anticipates relatively large risk and/or collects a lot of perception data. Then, her best explanation of the data minimizes Kullback-Leibler divergence from the true lottery among all lotteries, which yields the correct estimate.

Let $q^*(\mathbf{r}) = \sum_{i=1}^I p_i q_i^*(\mathbf{r})$ be the value of the lottery $\mathbf{q}^*(\mathbf{r})$. Proposition 6 implies:

Corollary 1. Consider a decision problem (\mathbf{r}, s) such that $q^*(\mathbf{r}) > [<] s$. Then, the probability that the DM chooses the lottery [the safe option] approaches 1 as $n \rightarrow \infty$.

To focus on the effect of the curvature of the encoding function, we set the sampling frequencies equal to the actual probabilities and compare the asymptotic estimated lottery value $q^*(\mathbf{r})$ with the true value r of the lottery \mathbf{r} .¹⁹

Proposition 7. Let the encoding function m be twice differentiable. Let $\pi_i = p_i$, and \mathbf{r} be a fixed lottery. The value of its Bayesian estimate almost surely converges to

$$r + \frac{1}{2} \frac{m''(r)}{m'(r)} \cdot \frac{1 + 4z(r)}{(1 + z(r))^2} \cdot \sigma^2(\mathbf{r}) + o(\sigma^2(\mathbf{r})), \quad (12)$$

as $n \rightarrow \infty$, where $z(r) = a\Delta m^{\prime 2}(r)$. The factor $\frac{1+4z(r)}{(1+z(r))^2}$ attains values in $(0, 4/3]$ and approaches 1 and 0 as $a\Delta \rightarrow 0$ and $a\Delta \rightarrow \infty$, respectively.

To interpret the result, recall that the risk premium of an expected-utility maximizer with Bernoulli utility u for a lottery \mathbf{r} with small risk is approximately $\frac{1}{2} \frac{u''(r)}{u'(r)} \sigma^2(\mathbf{r})$. The approximate risk premium of our DM is the same for $u(\cdot) = m(\cdot)$ but scaled by the positive factor $\frac{1+4z(r)}{(1+z(r))^2}$. The DM's bias in the valuation of the lottery relative to r arises because the DM deems risk a priori unlikely and therefore concludes that her perceived data are generated by a lottery with a smaller reward variance than the true variance. The underestimation of the variance leads to a mismatch to the perception data and this mismatch leads to a bias in the estimated mean of the lottery.

The dependence of the risk premium on the parameters Δ and a sheds light on two apparent instabilities of risk preferences pointed out by Rabin (2000) and Kahneman (2011). Kahnemann distinguishes between fast and slow modes of decision-making, where the fast mode favours the risk-attitudes found in prospect theory whereas the slow mode favours risk-neutrality.²⁰ If the amount of perception data collected by the DM increases with the time available for the decision, then time pressure is captured in our example by a low value of parameter a . In accord with Kahnemann, we find encoding-based risk attitudes when $a \rightarrow 0$. When our DM, who has anticipated little risk, encounters a risky lottery under time pressure, the relatively few data points that she has collected are best explained by

¹⁹We say that function $f(\mathbf{r})$ is $o(g(\mathbf{r}))$ if $f(\mathbf{r}_k)/g(\mathbf{r}_k) \rightarrow 0$ for any sequence \mathbf{r}_k such that $\sigma(\mathbf{r}_k) \rightarrow 0$. Specifically, a function is $o(\sigma^2)$ if it is negligible relative to σ^2 for lotteries with small σ . The expression $o(\cdot)$ stands for “term of smaller order than”.

²⁰Kirchler et al. (2017) show experimentally that time pressure increases risk aversion for gains and risk loving for losses. Relatedly, Porcelli and Delgado (2009) and Cahliková and Cingl (2017) find that stress accentuates risk attitudes in lab choices. But see also Kocher, Pahlke, and Trautmann (2013) who do not find an increase of risk aversion due to time pressure in their design.

an a priori likely low-risk lottery. Which such low-risk lottery is the best fit to the DM's data depends on the encoding function, thus the curvature of m determines the DM's risk attitudes. At the other extreme, in the absence of time pressure, when $a \rightarrow \infty$, the DM collects enough data for her prior to be irrelevant. She then learns the lottery and makes the risk-neutral choice.

Rabin (2000) points out that the risk-averse choices observed for small risks imply implausibly high risk aversion for large risks under a stable Bernoulli utility function. In our model, risk attitudes depend on the level of a priori anticipated risk. The anticipation of low risk – captured by small Δ here – induces risk attitudes since it makes risky lotteries surprising, and this leads to distortion of the posteriors when a risky lottery is encountered. If, however, the DM anticipates high-risk lotteries – if parameter Δ is large – then the DM's risk attitudes are attenuated. Risky lotteries become unsurprising and the DM's posterior expectation approaches the lottery's true expected value.

6 Summary

We develop a model of constrained optimal perception of gambles in which psychophysical adaptation affects choices. The impact of the perceptual strategy vanishes for rich perception data if the DM encounters a lottery that she has anticipated, but perception-induced risk attitudes arise for risk that the DM has not anticipated. In the latter case, we provide a unified explanation for various well-documented patterns in risky choice: adaptive risk attitudes, S-shaped reward valuation, probability weighting, and the role of stakes and time pressure.

The model makes several novel predictions. For example, explaining the structure of risk to the DM should attenuate her risk attitudes, while increasing the complexity of the environment should strengthen perception-driven behavior. These predictions are broadly in line with the recent experimental findings of Enke and Graeber (2021), who show that probability weighting is more pronounced for subjects who state a higher level of cognitive uncertainty about the correct action. Other predictions relate to the effect of framing. For example, a DM who is framed to perceive a risky lottery as riskless will rely on sampling frequencies rather than objective probabilities to evaluate the lottery. Manipulation of the sampling frequencies then has a strong impact on choice. A seller offering a risky prospect can make it more attractive if the presentation of the prospect leads to over-sampling of the upside risk. An additional prediction is that risk attitudes become more pronounced if the DM samples less perceptual data, which the seller of an insurance contract could exploit by putting the DM under time pressure.

References

- Attneave, F. (1954). Some informational aspects of visual perception. *Psychological review* 61(3), 183.
- Barlow, H. B. (1961). Possible principles underlying the transformation of sensory messages. *Sensory communication* 1(01).
- Benkert, J.-M. and N. Netzer (2018). Informational requirements of nudging. *Journal of Political Economy* 126(6), 2323–2355.
- Berk, R. H. (1966). Limiting behavior of posterior distributions when the model is incorrect. *The Annals of Mathematical Statistics*, 51–58.
- Bernheim, B. D. and A. Rangel (2009). Beyond revealed preference: choice-theoretic foundations for behavioral welfare economics. *The Quarterly Journal of Economics* 124(1), 51–104.
- Bethge, M., D. Rotermund, and K. Pawelzik (2002). Optimal short-term population coding: when fisher information fails. *Neural computation* 14(10), 2317–2351.
- Birnbaum, Z. W. (1948). On random variables with comparable peakedness. *The Annals of Mathematical Statistics* 19(1), 76–81.
- Cahlíková, J. and L. Cingl (2017). Risk preferences under acute stress. *Experimental Economics* 20(1), 209–236.
- Eliasz, K. and R. Spiegel (2020). A model of competing narratives. *American Economic Review* 110(12), 3786–3816.
- Enke, B. and T. Graeber (2021). Cognitive uncertainty. Mimeo.
- Esponda, I. and D. Pouzo (2016). Berk–nash equilibrium: A framework for modeling agents with misspecified models. *Econometrica* 84(3), 1093–1130.
- Fechner, G. T. (1860). Elements of psychophysics.
- Frick, M., R. Iijima, and Y. Ishii (2021). Welfare comparisons for biased learning.
- Friedman, D. (1989). The S-shaped value function as a constrained optimum. *American Economic Review* 79(5), 1243–1248.

- Frydman, C. and L. J. Jin (2019). Efficient coding and risky choice. *Available at SSRN 3270773*.
- Heidhues, P., B. Köszegi, and P. Strack (2018). Unrealistic expectations and misguided learning. *Econometrica* 86(4), 1159–1214.
- Heng, J. A., M. Woodford, and R. Polania (2020). Efficient sampling and noisy decisions. *Elife* 9, e54962.
- Herold, F. and N. Netzer (2015). Second-best probability weighting. Mimeo.
- Johnson, D. H. and G. C. Orsak (1993). Relation of signal set choice to the performance of optimal non-gaussian detectors. *IEEE Transactions on Communications* 41(9), 1319–1328.
- Juechems, K., J. Balaguer, B. Spitzer, and C. Summerfield (2021). Optimal utility and probability functions for agents with finite computational precision. *Proceedings of the National Academy of Sciences* 118(2).
- Kahneman, D. (2011). *Thinking, fast and slow*. Macmillan.
- Kahneman, D. and A. Tversky (1979). Prospect theory: An analysis of decision under risk. *Econometrica* 47(2), 263–291.
- Kahneman, D., P. P. Wakker, and R. Sarin (1997). Back to bentham? explorations of experienced utility. *The Quarterly Journal of Economics* 112(2), 375–406.
- Khaw, M. W., Z. Li, and M. Woodford (2018). Cognitive imprecision and small-stakes risk aversion. *The Review of Economic Studies*.
- Kirchler, M., D. Andersson, C. Bonn, M. Johannesson, E. Ø. Sørensen, M. Stefan, G. Tinghög, and D. Västfjäll (2017). The effect of fast and slow decisions on risk taking. *Journal of Risk and Uncertainty* 54(1), 37–59.
- Kocher, M. G., J. Pahlke, and S. T. Trautmann (2013). Tempus fugit: time pressure in risky decisions. *Management Science* 59(10), 2380–2391.
- Koszegi, B. and M. Rabin (2008). Revealed mistakes and revealed preferences. *The foundations of positive and normative economics: a handbook*, 193–209.
- Laughlin, S. (1981). A simple coding procedure enhances a neuron’s information capacity. *Zeitschrift für Naturforschung c* 36(9-10), 910–912.

- Le Cam, L. (1953). On some asymptotic properties of maximum likelihood estimates and related bayes estimates. *Univ. California Pub. Statist.* 1, 277–330.
- Lieder, F., T. Griffiths, and M. Hsu (2017). Overrepresentation of extreme events in decision making reflects rational use of cognitive resources. *Psychological Review* 125, 132.
- Molavi, P. (2019). Macroeconomics with learning and misspecification: A general theory and applications. *Unpublished manuscript*.
- Netzer, N. (2009). Evolution of time preferences and attitudes toward risk. *The American Economic Review* 99(3), 937–955.
- Ok, E. A. (2011). *Real analysis with economic applications*. Princeton University Press.
- Payzan-LeNestour, E. and M. Woodford (2021). Outlier blindness: A neurobiological foundation for neglect of financial risk. *Journal of Financial Economics* (in press).
- Porcelli, A. J. and M. R. Delgado (2009). Acute stress modulates risk taking in financial decision making. *Psychological Science* 20(3), 278–283.
- Purkayastha, S. (1998). Simple proofs of two results on convolutions of unimodal distributions. *Statistics & probability letters* 39(2), 97–100.
- Rabin, M. (2000). Risk aversion and expected-utility theory: A calibration theorem. *Econometrica* 68(5), 1281–1292.
- Rayo, L. and G. S. Becker (2007). Evolutionary efficiency and happiness. *Journal of Political Economy* 115(2), 302–337.
- Robson, A. J. (2001). The biological basis of economic behavior. *Journal of Economic Literature* 39(1), 11–33.
- Robson, A. J., L. A. Whitehead, and N. Robalino (2021). Adaptive hedonic utility. Technical report, Working paper.
- Salant, Y. and A. Rubinstein (2008). (a, f): choice with frames. *The Review of Economic Studies* 75(4), 1287–1296.
- Savage, L. J. (1954). *The Foundations of Statistics*. John Wiley & Sons.
- Schaffner, J., P. Tobler, T. Hare, and R. Polania (2021). Fitness-maximizing codes in early sensory areas. University of Zurich mimeo.

- Starmer, C. and R. Sugden (1993). Testing for juxtaposition and event-splitting effects. *Journal of risk and uncertainty* 6(3), 235–254.
- Steiner, J. and C. Stewart (2016). Perceiving prospects properly. *The American Economic Review* 106(7), 1601–31.
- Thurstone, L. L. (1927). A law of comparative judgment. *Psychological review* 34(4), 273.
- Vieider, F. (2021). Noisy coding and decisions under uncertainty. Mimeo.
- Wald, A. (1949). Note on the consistency of the maximum likelihood estimate. *The Annals of Mathematical Statistics* 20(4), 595–601.
- Wang, Z., A. A. Stocker, and D. D. Lee (2016). Efficient neural codes that minimize lp reconstruction error. *Neural computation* 28(12), 2656–2686.
- White, H. (1982). Maximum likelihood estimation of misspecified models. *Econometrica: Journal of the Econometric Society*, 1–25.
- Woodford, M. (2012). Prospect theory as efficient perceptual distortion. *American Economic Review* 102(3), 41–46.
- Woodford, M. (2020). Modeling imprecision in perception, valuation, and choice. *Annual Review of Economics* 12, 579–601.

A Asymptotic Loss Characterization

Let $0 < \underline{h} \leq \bar{h} < +\infty$ and $0 < \underline{m}' \leq \bar{m}' < +\infty$ be bounds on the functions h and m' . These bounds exist since the two functions are continuous on a compact interval.

Lemma 1. *Suppose that the encoding function is continuously differentiable and the reward density h is continuous. Let $r_J^* \in (\underline{r}, \bar{r})$ be a realization of the reward, and $q_{J,n}^B = \mathbb{E}[r_J \mid m_{J,n}]$ and $q_{J,n}^{ML} = m^{-1}(m_{J,n})$ its Bayesian and ML estimators. Then,*

$$(i) \quad \sqrt{n} (q_{J,n}^B - q_{J,n}^{ML}) \rightarrow 0 \text{ as } n \rightarrow \infty \text{ (a.s.)},$$

$$(ii) \quad n \text{Var}[r_J \mid m_{J,n}] \rightarrow \frac{1}{\pi_J m'^2(r_J^*)} \text{ as } n \rightarrow \infty \text{ (a.s.)},$$

$$(iii) \quad \text{The MSE of the maximum-likelihood estimate (rescaled by } n) \text{ is uniformly bounded:}$$

$$n \mathbb{E} \left[(r_J - q_{J,n}^{ML})^2 \mid m_{J,n} \right] \leq \frac{\bar{h}}{\underline{h}} \cdot \frac{\bar{m}'}{\underline{m}'^3} \frac{1}{\pi_J}.$$

Proof. Consider sufficiently large n so that $m_{J,n} \in [\underline{m}, \bar{m}]$. We introduce the rescaled error $\hat{\varepsilon}_{J,n} := \sqrt{\pi_J n}(r_J - q_{J,n})$ and derive its conditional density given $m_{J,n}$. Since $m_{J,n} \sim \mathcal{N}(m(r_J^*), 1/(\pi_J n))$, the pdf of $r_J \mid m_{J,n}$ is proportional to

$$h(\tilde{r}) \varphi\left(\sqrt{\pi_J n}(m(\tilde{r}) - m_{J,n})\right)$$

for any $\tilde{r} \in [\underline{r}, \bar{r}]$ and 0 otherwise; recall that φ is the standard normal density. Thus, the pdf of $\hat{\varepsilon}_{J,n}$ conditioned on $m_{J,n} = m(q_{J,n}^{ML})$ is

$$h_{J,n}(\tilde{\varepsilon}) = h_{J,n}^0 \cdot h\left(q_{J,n}^{ML} + \frac{\tilde{\varepsilon}}{\sqrt{\pi_J n}}\right) \varphi\left(\sqrt{\pi_J n}\left(m\left(q_{J,n}^{ML} + \frac{\tilde{\varepsilon}}{\sqrt{\pi_J n}}\right) - m(q_{J,n}^{ML})\right)\right)$$

for any $\tilde{\varepsilon} \in [\sqrt{\pi_J n}(\underline{r} - q_{J,n}^{ML}), \sqrt{\pi_J n}(\bar{r} - q_{J,n}^{ML})]$ and 0 otherwise; $h_{J,n}^0$ is the normalization factor. It follows that $h_{J,n}(\tilde{\varepsilon})/h_{J,n}^0$ is dominated by the integrable function $\bar{h} \cdot \varphi(\underline{m}' \cdot \tilde{\varepsilon})$.

Since $|q_{J,n}^{ML} - r_J^*| = |m^{-1}(m_{J,n}) - r_J^*| \leq \frac{1}{\underline{m}'} |m_{J,n} - m(r_J^*)|$ and $m_{J,n} \sim \mathcal{N}(m(r_J^*), 1/(\pi_J n))$, we have that $q_{J,n}^{ML} \rightarrow r_J^*$ (a.s.). Using this, the Mean Value Theorem, and continuity of m' , we get

$$\sqrt{\pi_J n}\left(m\left(q_{J,n}^{ML} + \frac{\tilde{\varepsilon}}{\sqrt{\pi_J n}}\right) - m(q_{J,n}^{ML})\right) \rightarrow m'(r_J^*) \tilde{\varepsilon} \text{ as } n \rightarrow \infty \text{ (a.s.)},$$

and thus, using the continuity of h , for any $\tilde{\varepsilon}$,

$$\frac{h_{J,n}(\tilde{\varepsilon})}{h_{J,n}^0} \rightarrow h(r_J^*) \varphi(m'(r_J^*) \tilde{\varepsilon}) \text{ as } n \rightarrow \infty \text{ (a.s.)}.$$

Next, we characterize the limit of the normalization factors. By the Dominated Convergence Theorem,

$$\int_{\mathbb{R}} \frac{h_{J,n}(\tilde{\varepsilon})}{h_{J,n}^0} d\tilde{\varepsilon} \rightarrow \frac{1}{h_J^0} \text{ as } n \rightarrow \infty \text{ (a.s.)}, \text{ where } h_J^0 := \left[\int_{\mathbb{R}} h(r_J^*) \varphi(m'(r_J^*) \tilde{\varepsilon}) d\tilde{\varepsilon} \right]^{-1}.$$

Since $\int_{\mathbb{R}} h_{J,n}(\tilde{\varepsilon}) d\tilde{\varepsilon} = 1$ for all n , it follows that $h_{J,n}^0 \rightarrow h_J^0 > 0$ (a.s.). In particular, $h_{J,n}^0$ is bounded. Then, the posterior errors $\hat{\varepsilon}_{J,n} \mid m_{J,n}$ converge in distribution to $\mathcal{N}(0, 1/m'^2(r_J^*))$ (a.s.).

Applying the Dominated Convergence Theorem to the functions $\tilde{\varepsilon} h_{J,n}(\tilde{\varepsilon})$ and $\tilde{\varepsilon}^2 h_{J,n}(\tilde{\varepsilon})$, we conclude that $E[\hat{\varepsilon}_{J,n} \mid m_{J,n}] \rightarrow 0$ and $\text{Var}[\hat{\varepsilon}_{J,n} \mid m_{J,n}] \rightarrow 1/m'^2(r_J^*)$ as $n \rightarrow \infty$ (a.s.). Claims (i) and (ii) follow from deriving r_J from $\hat{\varepsilon}_{J,n} = \sqrt{\pi_J n}(r_J - q_{J,n}^{ML})$.

For Claim (iii), recall that $h_{J,n}(\tilde{\varepsilon})/h_{J,n}^0$ is dominated by the integrable function $\bar{h} \cdot \varphi(\underline{m}' \cdot \tilde{\varepsilon})$ that equals, up to a multiplicative constant, the pdf of $\mathcal{N}(0, 1/\underline{m}'^2)$. Consider a random variable $\hat{\varepsilon}'_{J,n}$ with pdf proportional to $h_{J,n}(\tilde{\varepsilon})/h_{J,n}^0$ on the domain of $\hat{\varepsilon}_{J,n}$, and $\underline{h} \varphi(\bar{m}' \cdot \tilde{\varepsilon})$

outside of the domain. We can establish the following upper bound on the normalization constant $h_{J,n}^0$ of the pdf of the variable $\varepsilon'_{J,n}$,

$$h_{J,n}^0 \leq \left[\int_{\mathbb{R}} \underline{h} \varphi(\overline{m}' \cdot \tilde{\varepsilon}) d\tilde{\varepsilon} \right]^{-1} = \frac{\overline{h}}{\underline{h}} \cdot \frac{\overline{m}'}{\underline{m}'} \cdot \left[\int_{\mathbb{R}} \overline{h} \varphi(\underline{m}' \cdot \tilde{\varepsilon}) d\tilde{\varepsilon} \right]^{-1}.$$

Then,

$$n \mathbb{E} \left[(r_J - q_{J,n}^{ML})^2 \mid m_{J,n} \right] \leq \frac{1}{\pi_J} \mathbb{E} \left[\varepsilon'^2_{J,n} \mid m_{J,n} \right] \leq \frac{1}{\pi_J} h_{J,n}^0 \int_{\mathbb{R}} \tilde{\varepsilon}^2 \cdot \overline{h} \varphi(\underline{m}' \tilde{\varepsilon}) d\tilde{\varepsilon} = \frac{\overline{h}}{\underline{h}} \cdot \frac{\overline{m}'}{\underline{m}'^3} \frac{1}{\pi_J}.$$

□

Corollary 2. *Conditional on a realization of $\mathbf{r}^* \in (\underline{r}, \overline{r})^{|\mathcal{J}|}$,*

$$(i) \sqrt{n} (q_n^B - q_n^{ML}) \rightarrow 0 \text{ as } n \rightarrow \infty \text{ (a.s.)},$$

$$(ii) n \text{Var}[r \mid \mathbf{m}_n] \rightarrow \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J^*)} \text{ as } n \rightarrow \infty \text{ (a.s.)}.$$

$$(iii) n \mathbb{E} \left[(r - q_n^{ML})^2 \mid \mathbf{m}_n \right] \leq \frac{\overline{h}}{\underline{h}} \cdot \frac{\overline{m}'}{\underline{m}'^3} \cdot \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J}.$$

For the following two lemmas, we abstract from the specific structure of the messages \mathbf{m}_n , let the DM receive a vector of messages $\mathbf{m} = (m_J)_J$ and then form the estimate $q(\mathbf{m})$ of r as a function of \mathbf{m} . Let

$$\ell := \max\{r, s\} - \mathbb{1}_{q > s} r - \mathbb{1}_{q \leq s} s$$

denote the loss from using the estimate q .

We assume that the pdf h_s is continuously differentiable, thus h_s and h'_s are bounded from above; let \overline{h}_s and \overline{h}'_s be the respective bounds. We say that $O(\cdot)$ has uniform bound \overline{h}'_s if $|O(x)/x| \leq \overline{h}'_s$ for all x and any value of \mathbf{r} and \mathbf{m} .²¹

Lemma 2. *The expected loss of the estimate q conditioned on \mathbf{m} and \mathbf{r} is*

$$\mathbb{E}[\ell \mid \mathbf{r}, \mathbf{m}] = \frac{1}{2} h_s(q) (r - q)^2 + O((r - q)^3),$$

where the expectation is over s and $O(\cdot)$ has the uniform bound \overline{h}'_s .

Proof. Consider a fixed realization of \mathbf{r} and \mathbf{m} . The loss is $\ell = |r - s|$ if the DM makes the suboptimal choice, which happens if and only if s is between the true lottery value r and its

²¹The term $O(\cdot)$ stands for the “term of the order of”.

estimate q . Taking the expectation over the safe option yields (for both $r < q$ and $r > q$)

$$\mathbb{E}[\ell \mid \mathbf{r}, \mathbf{m}] = \int_r^q (\tilde{s} - r) h_s(\tilde{s}) d\tilde{s}.$$

The lemma follows from the approximation $h_s(\tilde{s}) = h_s(q) + O(\tilde{s} - q)$, in which $O(\cdot)$ has the uniform bound \bar{h}'_s ,

$$\begin{aligned} \int_r^q (\tilde{s} - r) h_s(\tilde{s}) d\tilde{s} &= h_s(q) \int_r^q (\tilde{s} - r) d\tilde{s} + \int_r^q (\tilde{s} - r) O(\tilde{s} - q) d\tilde{s} \\ &= \frac{1}{2} h_s(q) (r - q)^2 + O((r - q)^3). \end{aligned}$$

□

Lemma 3. *The expected loss of the estimate q conditioned on \mathbf{m} is*

$$\mathbb{E}[\ell \mid \mathbf{m}] = \frac{1}{2} h_s(q) \sigma^2 + O(\sigma^3), \text{ where } \sigma^2 := \text{Var}[r \mid \mathbf{m}] + (q^B - q)^2,$$

where $O(\cdot)$ has the uniform bound \bar{h}'_s and $q^B = \mathbb{E}[r \mid \mathbf{m}]$.

Proof. This follows from Lemma 2 by taking the expectation over \mathbf{r} :

$$\begin{aligned} \mathbb{E}[\ell \mid \mathbf{m}] &= \mathbb{E} \left[\mathbb{E}[\ell \mid \mathbf{r}, \mathbf{m}] \mid \mathbf{m} \right] \\ &= \mathbb{E} \left[\frac{1}{2} h_s(q) (r - q)^2 + O((r - q)^3) \mid \mathbf{m} \right], \end{aligned}$$

where $O(\cdot)$ has the uniform bound \bar{h}'_s . Since $|O((r - q)^3)| \leq \bar{h}'_s |r - q|^3$,

$$\left| \mathbb{E} [O((r - q)^3) \mid \mathbf{m}] \right| \leq \mathbb{E} \left[\bar{h}'_s \left((r - q)^2 \right)^{3/2} \mid \mathbf{m} \right] \leq \bar{h}'_s \mathbb{E} [(r - q)^2 \mid \mathbf{m}]^{3/2},$$

where we have used Jensen's inequality in the second step.

We conclude with

$$\begin{aligned} \mathbb{E} [(r - q)^2 \mid \mathbf{m}] &= \mathbb{E} \left[\left((r - q^B) + (q^B - q) \right)^2 \mid \mathbf{m} \right] \\ &= \mathbb{E} \left[(r - q^B)^2 + 2(r - q^B)(q^B - q) + (q^B - q)^2 \mid \mathbf{m} \right] \\ &= \text{Var}[r \mid \mathbf{m}] + (q^B - q)^2 = \sigma^2. \end{aligned}$$

□

Proof of Proposition 1. Let $\ell_n^z = \max\{r, s\} - \mathbb{1}_{q_n^z > s} r - \mathbb{1}_{q_n^z \leq s} s$ be the loss of the estimator

q_n^z , $z \in \{B, ML\}$. For a given realization of the lottery \mathbf{r}^* with value r^* , we prove that the expected loss conditioned on \mathbf{m}_n satisfies

$$n \mathbb{E}[\ell_n^z \mid \mathbf{m}_n] \rightarrow \frac{1}{2} h_s(r^*) \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J^*)} \text{ as } n \rightarrow \infty \text{ (a.s.)}, \quad (13)$$

where the expectation is over s and q_n^z is a function of \mathbf{m}_n .

Lemma 3 applied to \mathbf{m}_n and q_n^z implies

$$n \mathbb{E}[\ell_n^z \mid \mathbf{m}_n] = \frac{1}{2} h_s(q_n^z) n \sigma_n^2(z) + n O(\sigma_n^3(z)), \text{ where } \sigma_n^2(z) := \text{Var}[r \mid \mathbf{m}_n] + (q_n^B - q_n^z)^2. \quad (14)$$

Corollary 2 implies that $n(q_n^B - q_n^z)^2 \rightarrow 0$ (a.s.) (this holds trivially for the Bayesian estimator). Further, Claim (ii) of Corollary 2 implies

$$n \sigma_n^2(z) \rightarrow \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J^*)} \text{ as } n \rightarrow \infty \text{ (a.s.)}.$$

Thus, $\sigma_n(z) \rightarrow 0$ as $n \rightarrow \infty$ (a.s.); and so $|n O(\sigma_n^3(z))| \leq n \bar{h}'_s \sigma_n^3(z) = \bar{h}'_s \cdot n \sigma_n^2(z) \cdot \sigma_n(z) \rightarrow 0$ as $n \rightarrow \infty$ (a.s.). Substituting back into (14) and taking into account that $q_n^z \rightarrow r^*$ as $n \rightarrow \infty$ (a.s.), we obtain (13).

The proposition follows from taking expectation over \mathbf{r} and applying the Dominated Convergence Theorem. In particular, (14) implies that $n \mathbb{E}[\ell_n^z \mid \mathbf{m}_n]$ has an integrable bound:

$$|n \mathbb{E}[\ell_n^z \mid \mathbf{m}_n]| \leq \frac{1}{2} \bar{h}_s \cdot n \sigma_n^2(ML) + \bar{h}'_s \cdot n \sigma_n^2(ML) \cdot \sigma_n(ML) \leq \frac{1}{2} \bar{h}_s \bar{\Sigma} + \bar{h}'_s \bar{\Sigma} \cdot \max\{\bar{\Sigma}, 1\},$$

where $\bar{\Sigma} = \frac{\bar{h}}{h} \cdot \frac{\bar{m}'}{m'^3} \cdot \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J}$ is the uniform bound from Claim (iii) of Corollary 2. \square

B Optimal Perception

B.1 Proof of Proposition 2

Proof of Proposition 2. The objective of the information-processing problem is a functional

$$\mathcal{L}(m'(\cdot), (\pi_J)_{J \in \mathcal{J}}) = \mathbb{E} \left[\sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J)} \mid r = s \right].$$

Since $\frac{p_J^2}{\pi_J m'^2(r_J)}$ is convex with respect to $(m'(r_J), \pi_J)$, the functional \mathcal{L} is convex. Thus, the first-order conditions are sufficient for a global minimum of the information-processing

problem.

Since the objective (4) is strictly decreasing in both $m'(\cdot)$ and π_J , the constraints (5) and (6) are binding at the optimum. The Lagrangian of the constrained optimization problem (4)-(6) is

$$\begin{aligned} & \mathbb{E} \left[\sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J)} \mid r = s \right] + \lambda \left(\int_{\underline{r}}^{\bar{r}} m'(\tilde{r}) d\tilde{r} - (\bar{m} - \underline{m}) \right) + \mu \left(\sum_{J \in \mathcal{J}} \pi_J - 1 \right) = \\ & \sum_{J \in \mathcal{J}} \int_{\underline{r}}^{\bar{r}} \frac{p_J^2}{\pi_J m'^2(\tilde{r}_J)} h_J(\tilde{r}_J) d\tilde{r}_J + \lambda \left(\int_{\underline{r}}^{\bar{r}} m'(\tilde{r}) d\tilde{r} - (\bar{m} - \underline{m}) \right) + \mu \left(\sum_{J \in \mathcal{J}} \pi_J - 1 \right), \end{aligned}$$

where λ and μ are the Lagrange multipliers for (5) and (6), respectively. The first-order condition (9) then follows by summing the derivatives w.r.t. $m'(\tilde{r})$, $\tilde{r} \in [\underline{r}, \bar{r}]$, of all the integrands in the last inline expression. This first-order condition must hold for almost all \tilde{r} , so the optimal encoding function m satisfies the condition for all \tilde{r} . Expressing $m'(\tilde{r})$ from (9) gives (7). Further, m' is continuous since each h_J is continuous.

The first-order condition of the information-processing problem with respect to π_J is, for each $J \in \mathcal{J}$,

$$\left(\frac{p_J}{\pi_J} \right)^2 \mathbb{E} \left[\frac{1}{m'^2(r_J)} \mid r = s \right] = \mu.$$

This implies (8). □

B.2 Proof of Proposition 3

Definition 1. *A continuous random variable is unimodal and symmetric around 0 if its density function $h(x)$ is strictly decreasing on the positive part of its domain and $h(x) = h(-x)$ for all $x \in \mathbb{R}$.*

This property is preserved by summation: the sum of unimodal and symmetric random variables is unimodal and symmetric, see e.g. Purkayastha (1998).

Definition 2 (Birnbbaum (1948)). *Let X and Y be two unimodal random variables symmetric around 0. We say that X is more peaked than Y if $P(|X| < \alpha) > P(|Y| < \alpha)$ (unless the right-hand side is 1) for all $\alpha > 0$.*

Equivalently, for two unimodal symmetric random variables, X is more peaked than Y whenever the cdf of X is greater than the cdf of Y at any $\alpha > 0$ from the support of Y .

For the next two lemmas, let X_0, X_1, \dots, X_I be independent real-valued continuous random variables that are unimodal and symmetric around 0, where X_1, \dots, X_I are identically

distributed and the distribution of X_0 may be distinct from that of X_i , $i > 0$. Denote by h the pdf of each of the iid variables X_1, \dots, X_I . Let $(p_1, \dots, p_I) \in \Delta(\{1, \dots, I\})$, and $X := \sum_{i=1}^I p_i X_i$.

Lemma 4. *The random variable $X_i \mid (X = X_0)$, $i = 1, \dots, I$, is unimodal and symmetric around 0.*

Proof. Since unimodality together with symmetry is preserved by affine combinations, the variable $X_{-i} := \frac{1}{p_i}(X_0 - \sum_{k \neq i} p_k X_k)$ is unimodal and symmetric around 0. Denote by h_{-i} the pdf of X_{-i} . Then $X_i \mid (X = X_0)$ is identical to $X_i \mid (X_i = X_{-i})$, and so its pdf is, up to a normalization constant, $h(x_i)h_{-i}(x_i)$, which is unimodal and symmetric around 0, as those properties are preserved when taking product of pdfs. \square

Lemma 5. *The random variable $X_i \mid (X = X_0)$ is more peaked than $X_j \mid (X = X_0)$ if and only if $p_i > p_j$.*

Proof. Without loss of generality, assume $\{i, j\} = \{1, 2\}$ (that is, either $i = 1$ and $j = 2$ or $i = 2$ and $j = 1$). Define $X_{-12} := X_0 - \sum_{k=3}^I p_k X_k$ (if $I = 2$, then $X_{-12} = X_0$) and let h_{-12} be its pdf. This is a unimodal random variable symmetric around 0. The random variable $X_i \mid (X = X_0)$ is identical to $X_i \mid (p_i X_i + p_j X_j = X_{-12})$ and so its pdf equals

$$h_i(x_i) = \frac{\int_{\mathbb{R}} h_{-12}(p_1 x_1 + p_2 x_2) h(x_1) h(x_2) dx_j}{\mathbb{E}[h_{-12}(p_1 X_1 + p_2 X_2)]},$$

where the expectation, which is with respect to X_1 and X_2 , is independent of i . Thus, for any $\alpha > 0$,

$$P(|X_1| < \alpha \mid X = X_0) = \frac{\iint_{(-\alpha, \alpha) \times \mathbb{R}} h_{-12}(p_1 x_1 + p_2 x_2) h(x_1) h(x_2) dx_1 dx_2}{\mathbb{E}[h_{-12}(p_1 X_1 + p_2 X_2)]}$$

$$P(|X_2| < \alpha \mid X = X_0) = \frac{\iint_{(-\alpha, \alpha) \times \mathbb{R}} h_{-12}(p_1 x_2 + p_2 x_1) h(x_1) h(x_2) dx_1 dx_2}{\mathbb{E}[h_{-12}(p_1 X_1 + p_2 X_2)]},$$

where we used that $P(|X_1| < \alpha \mid X = X_0)$ and $P(|X_2| < \alpha \mid X = X_0)$ are both (up to the same normalization constant) integrals of the same function $(x_1, x_2) \mapsto h_{-12}(p_1 x_1 + p_2 x_2) h(x_1) h(x_2)$, but the first is over the region $[-\alpha, \alpha] \times \mathbb{R}$, and the second is over $\mathbb{R} \times [-\alpha, \alpha]$. This is equivalent to integrating both over the same region but switching the roles of x_1 and x_2 . Then,

$$(P(|X_1| < \alpha \mid X = X_0) - P(|X_2| < \alpha \mid X = X_0)) \cdot \mathbb{E}[h_{-12}(p_1 X_1 + p_2 X_2)] =$$

$$\begin{aligned}
& \iint_{(-\alpha, \alpha) \times \mathbb{R}} \left(h_{-12}(p_1 x_1 + p_2 x_2) - h_{-12}(p_1 x_2 + p_2 x_1) \right) h(x_1) h(x_2) dx_1 dx_2 = \\
& \iint_{(-\alpha, \alpha) \times (\mathbb{R} \setminus (-\alpha, \alpha))} \left(h_{-12}(p_1 x_1 + p_2 x_2) - h_{-12}(p_1 x_2 + p_2 x_1) \right) h(x_1) h(x_2) dx_1 dx_2 = \\
& 2 \iint_{(-\alpha, \alpha) \times [\alpha, +\infty)} \left(h_{-12}(p_1 x_1 + p_2 x_2) - h_{-12}(p_1 x_2 + p_2 x_1) \right) h(x_1) h(x_2) dx_1 dx_2,
\end{aligned}$$

where we used that both integrals cancel each other out on the region $(-\alpha, \alpha) \times (-\alpha, \alpha)$, and that h and h_{-12} are symmetric around 0.

Suppose that $p_2 > p_1$, and consider any $(x_1, x_2) \in (-\alpha, \alpha) \times [\alpha, +\infty)$. It follows from the identity

$$p_1 x_1 + p_2 x_2 = (p_1 x_2 + p_2 x_1) + (p_2 - p_1)(x_2 - x_1)$$

that

$$p_1 x_1 + p_2 x_2 > p_1 x_2 + p_2 x_1,$$

where the left-hand side (LHS) is always positive. The right-hand side (RHS) is either positive or negative, but smaller in absolute value than the LHS. Indeed, if the RHS is negative, then $x_1 < 0$, and

$$|p_1 x_2 + p_2 x_1| = -p_1 x_2 + p_2 |x_1| = -p_1 |x_1| + p_2 x_2 - (p_1 + p_2)(x_2 - |x_1|) < -p_1 |x_1| + p_2 x_2.$$

Thus,

$$|p_1 x_1 + p_2 x_2| > |p_1 x_2 + p_2 x_1|,$$

and due to the symmetry and unimodality of h_{-12} ,

$$h_{-12}(p_1 x_1 + p_2 x_2) < h_{-12}(p_1 x_2 + p_2 x_1),$$

unless both are zero. It follows that $X_2 \mid (X = X_0)$ is more peaked than $X_1 \mid (X = X_0)$, as needed. \square

Lemma 6. *Let the function f be continuous, symmetric around 0 and increasing on \mathbb{R}_+ , and let X_1, X_2 be unimodal continuous random variables that are symmetric around 0 and have bounded support. Then $\mathbb{E}[f(X_1)] < \mathbb{E}[f(X_2)]$ whenever X_1 is more peaked than X_2 .*

Proof. Denote by $h_i(x)$ and $H_i(x)$ the pdf and cdf of X_i , $i = 1, 2$. Then,

$$\frac{1}{2} \mathbb{E}[f(X_i)] = \int_0^\infty f(x) h_i(x) dx$$

$$\begin{aligned}
&= \left[f(x)(H_i(x) - 1) \right]_0^{+\infty} - \int_0^{+\infty} (H_i(x) - 1)df(x) \\
&= \frac{1}{2}f(0) + \int_0^{+\infty} (1 - H_i(x))df(x),
\end{aligned}$$

where we have used integration by parts for the Stieltjes integral, see e.g. Ok (2011). If X_1 is more peaked than X_2 , then $1 - H_1(x) < 1 - H_2(x)$ unless both are zero for all $x > 0$. It follows that $E[f(X_1)] < E[f(X_2)]$. \square

Proof of Proposition 3. Statement 1 follows from (7) because by Lemma 4 each h_J is unimodal with the same mode as the unconditional reward density h . Additionally, m' is symmetric around r_m since each h_J is symmetric around r_m .

Now consider Statement 2. Suppose $p_J < p_{J'}$. By (8) it suffices to show that

$$E \left[\frac{1}{m'^2(r_J)} \mid r = s \right] > E \left[\frac{1}{m'^2(r_{J'})} \mid r = s \right]. \quad (15)$$

This indeed holds since, by Lemma 5, $r_{J'} \mid (r = s)$ is more peaked than $r_J \mid (r = s)$ and the inequality (15) follows from Lemma 6 and from the fact that $1/m'^2(r)$ is continuous and symmetric around r_m and increasing above r_m . \square

B.3 Extension

We discuss here an extension of Proposition 2 and Proposition 3 to a setting in which the DM does not know the payoff-relevant partition of states \mathcal{J} at the point of optimization of the perception strategy, but knows the distribution of possible partitions. The timing is as follows: first, the DM chooses her perception strategy, i.e., the encoding function m and sampling frequencies π_i for all states $i = 1, \dots, I$. Afterwards, \mathcal{J} is realized and observed by the DM. The DM then samples the realized vector $\mathbf{r} \in A_{\mathcal{J}}$ according to her perception strategy. As in Section 4, the rewards are iid from a density h and the safe option s is independently drawn from a density h_s (both independent of \mathcal{J}). The DM observes s , forms an estimate of the lottery value r using her knowledge of the partition \mathcal{J} , and makes the optimal choice.

Proposition 1 applies for each realization of the partition. Hence, the limit loss (rescaled by $2n$) equals

$$\frac{1}{2} E \left[h_s(r) \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^2(r_J)} \right], \quad (16)$$

where the expectation is with respect to the partition \mathcal{J} and the reward vector \mathbf{r} . The objec-

tive probabilities and effective sampling frequencies for each element J of a given partition \mathcal{J} are $p_J = \sum_{i \in J} p_i$ and $\pi_J = \sum_{i \in J} \pi_i$, respectively. The information-processing problem for this setting is to minimize (16) subject to constraint (5) from the main text and $\sum_{i=1}^I \pi_i = 1$, which replaces the constraint (6).

The first-order condition for the slope $m'(\tilde{r})$ of the encoding function is

$$\mathbb{E} \left[2 \sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J m'^3(\tilde{r})} h_J(\tilde{r}) \right] = \lambda$$

for each \tilde{r} , where λ is the shadow price of the constraint (5) and the expectation is with respect to the partition. From this we obtain

$$m'(\tilde{r}) \propto \mathbb{E} \left[\sum_{J \in \mathcal{J}} \frac{p_J^2}{\pi_J} h_J(\tilde{r}) \right]^{\frac{1}{3}},$$

which generalizes Statement 1 of Proposition 2. If the densities h and h_s are symmetric and unimodal with a same mode, then each conditional reward density h_J for each possible partition is symmetric and unimodal. Thus, the optimal encoding function is again S-shaped, which generalizes Statement 1 of Proposition 3.

The first-order condition with respect to π_i is, for each $i = 1, \dots, I$,

$$\mathbb{E} \left[\frac{p_{J(i)}^2}{\pi_{J(i)}^2 m'^2(r_{J(i)})} \mid r = s \right] = \mu,$$

where the expectation is with respect to the partition and the reward vector, $J(i)$ is the element of the realized partition that contains i , and μ is the shadow price of the sampling constraint $\sum_{i=1}^I \pi_i = 1$. Hence, for all $i, i' = 1, \dots, I$,

$$\mathbb{E} \left[\frac{p_{J(i)}^2}{\pi_{J(i)}^2 m'^2(r_{J(i)})} \mid r = s \right] = \mathbb{E} \left[\frac{p_{J(i')}^2}{\pi_{J(i')}^2 m'^2(r_{J(i')})} \mid r = s \right], \quad (17)$$

which generalizes Statement 2 of Proposition 2.

If m was linear, the solution of the first-order condition would be proportional sampling $\pi_i = p_i$. If m is S-shaped, qualitatively, states i which are often included in low-probability events should be oversampled, because high-probability events have more peaked reward distributions conditional on ties, and the S-shaped encoding function thus measures low-probability rewards less precisely in expectation. This intuition generalizes Statement 2 of Proposition 3. For illustration, assume that only two partitions arise with positive proba-

bility, the coarsest one $\mathcal{J} = \{\{1, \dots, I\}\}$ and the finest one $\mathcal{J}' = \{\{1\}, \dots, \{I\}\}$. Then (17) reduces to (8) for the finest partition, and we obtain oversampling of low-probability states exactly like in Proposition 3.

C Proofs of Propositions 4 and 5

Proposition 4 follows from Proposition 5 for $\mathcal{K} = \{\{1, \dots, I\}\}$.

Proof of Proposition 5. Let $f_{\mathbf{r}}(x)$ be the signal density conditional on the encountered lottery \mathbf{r} . That is, for signal $x = (\hat{m}, i)$, $f_{\mathbf{r}}(x) = \pi_i \varphi(\hat{m} - m(r_i))$ where φ is the standard normal density. Kullback-Leibler divergence of the signal densities for any two lotteries \mathbf{r}, \mathbf{r}' is

$$\begin{aligned}
D_{\text{KL}}(f_{\mathbf{r}} \parallel f_{\mathbf{r}'}) &= \int_{\mathbb{R} \times \{1, \dots, I\}} f_{\mathbf{r}}(x) \ln \frac{f_{\mathbf{r}}(x)}{f_{\mathbf{r}'}(x)} dx \\
&= \sum_{i=1}^I \int_{\mathbb{R}} \pi_i \varphi(\hat{m} - m(r_i)) \ln \frac{\pi_i \varphi(\hat{m} - m(r_i))}{\pi_i \varphi(\hat{m} - m(r'_i))} d\hat{m} \\
&= \sum_{i=1}^I \pi_i \int_{\mathbb{R}} \varphi(\hat{m} - m(r_i)) \ln \frac{\varphi(\hat{m} - m(r_i))}{\varphi(\hat{m} - m(r'_i))} d\hat{m} \\
&= \sum_{i=1}^I \pi_i D_{\text{KL}}(\varphi_{m(r_i)} \parallel \varphi_{m(r'_i)}) \\
&= \frac{1}{2} \sum_{i=1}^I \pi_i (m(r_i) - m(r'_i))^2.
\end{aligned}$$

where $\varphi_m(\hat{m}) = \varphi(\hat{m} - m)$ is the density of the perturbed message \hat{m} conditional on the unperturbed message m . The last equality follows from the fact that the Kullback-Leibler divergence of two Gaussian densities with means μ_1, μ_2 and variances equal to 1 is $(\mu_1 - \mu_2)^2 / 2$ (see e.g. Johnson and Orsak, 1993).

Let

$$\mathbf{q} = \arg \min_{\mathbf{r}' \in \mathcal{A}_{\mathcal{K}}} D_{\text{KL}}(f_{\mathbf{r}} \parallel f_{\mathbf{r}'}) = \arg \min_{\mathbf{r}' \in \mathcal{A}_{\mathcal{K}}} \sum_{i=1}^I \pi_i (m(r_i) - m(r'_i))^2.$$

This minimizer $\mathbf{q} = (q_i)_i$ is unique and satisfies for each state $i = 1, \dots, I$,

$$m(q_i) = \arg \min_{m \in [\underline{m}, \bar{m}]} \sum_{j \in J(i)} \pi_j (m(r_j) - m)^2$$

$$= \sum_{j \in J(i)} \frac{\pi_j}{\pi_{J(i)}} m(r_j),$$

where $J(i)$ is the element of the partition \mathcal{K} that contains i .

The estimated lottery value q_n^z , $z \in \{ML, B\}$, almost surely converges to $\sum_{i=1}^I p_i q_i$. For the maximum-likelihood estimate, this follows from White (1982) who proves that it almost surely converges to the minimizer of the Kullback-Leibler divergence (provided the minimizer is unique). For the Bayesian estimate, the result follows from Berk (1966) who proves that the posterior belief almost surely converges in probability to an atom on the minimizer of the Kullback-Leibler divergence (again, provided the minimizer is unique). \square

D Proofs for Subsection 5.3

We use the next lemma in the proof of Proposition 6.

Lemma 7. *Let $\psi_n(\mathbf{x}) : [\underline{r}, \bar{r}]^I \rightarrow \mathbb{R}$ be a sequence of continuous functions uniformly converging to a function $\psi(\mathbf{x})$ which has a unique minimizer \mathbf{x}^* . Then, the random variable X_n with pdf equal to $\alpha_n \exp(-n\psi_n(\mathbf{x}))$, where α_n is the normalization factor, converges to \mathbf{x}^* in probability as $n \rightarrow \infty$.*

Proof. We need to prove that for every $\delta > 0$, the probability $P(X_n \in B_\delta) \rightarrow 1$ as $n \rightarrow \infty$, where B_δ is the open Euclidean δ -ball centered at \mathbf{x}^* . Fix $\delta > 0$ and define

$$d = \min_{\mathbf{x} \in [\underline{r}, \bar{r}]^I \setminus B_\delta} \{\psi(\mathbf{x}) - \psi(\mathbf{x}^*)\}.$$

The minimum exists as ψ is continuous and the set $[\underline{r}, \bar{r}]^I \setminus B_\delta$ is closed. Additionally, $d > 0$ since \mathbf{x}^* is the unique minimizer of ψ on $[\underline{r}, \bar{r}]^I$.

Because the convergence $\psi_n \rightarrow \psi$ is uniform, for any $d' > 0$ there exists $n_{d'} \in \mathbb{N}$ such that $|\psi_n(\mathbf{x}) - \psi(\mathbf{x})| < d'$ for all $\mathbf{x} \in [\underline{r}, \bar{r}]^I$ and $n \geq n_{d'}$. Consider $n \geq n_{d/4}$. Because $\psi_n(\mathbf{x}) \geq \psi(\mathbf{x}) - \frac{d}{4} \geq \psi(\mathbf{x}^*) + \frac{3d}{4}$ for \mathbf{x} outside of the ball B_δ , the probability density of X_n is at most $\alpha_n \exp(-n\psi(\mathbf{x}^*) - \frac{3d}{4}n)$. This implies,

$$P(X_n \notin B_\delta) \leq \tilde{\alpha}_n \exp\left(-\frac{3d}{4}n\right) (\bar{r} - \underline{r})^I, \quad \text{where } \tilde{\alpha}_n := \alpha_n \exp(-n\psi(\mathbf{x}^*)). \quad (18)$$

We conclude by establishing an upper bound for $\tilde{\alpha}_n$. Given $\delta > 0$, let $\delta' > 0$ be such that $\psi(\mathbf{x}) \leq \psi(\mathbf{x}^*) + d/4$ for all $\mathbf{x} \in B_{\delta'} \cap [\underline{r}, \bar{r}]^I$. Existence of such δ' follows from the continuity of ψ . Then, $\psi_n(\mathbf{x}) \leq \psi(\mathbf{x}) + \frac{d}{4} \leq \psi(\mathbf{x}^*) + \frac{d}{2}$ for all $\mathbf{x} \in B_{\delta'} \cap [\underline{r}, \bar{r}]^I$ and $n > n_{d/4}$. Thus the

probability density of X_n is at least $\tilde{\alpha}_n \exp(-\frac{d}{2}n)$ on this set. It follows that,

$$1 \geq P(X_n \in B_{\delta'}) \geq \tilde{\alpha}_n \exp\left(-\frac{d}{2}n\right) b',$$

where $b' > 0$ is the volume of the set $B_{\delta'} \cap [\underline{r}, \bar{r}]^I$. Substituting the implied upper bound on $\tilde{\alpha}_n$ into (18) gives

$$P(X_n \notin B_{\delta}) \leq \exp\left(-\frac{d}{4}n\right) \frac{(\bar{r} - \underline{r})^I}{b'}.$$

Since the right-hand vanishes as $n \rightarrow \infty$, the claim follows. \square

Proof of Proposition 6. Let $\mathbf{m}_n = (m_{i,n})_{i=1}^I$ be the vector of the averages of $a\pi n$ perturbed messages received for each state i . Since the encoding errors are standard normal, $m_{i,n} \mid r_i \sim \mathcal{N}\left(m(r_i), \frac{1}{a\pi_i n}\right)$. By Bayes' Rule, the posterior density of each lottery $\mathbf{r}' \in [\underline{r}, \bar{r}]^I$, is for given \mathbf{m}_n , proportional to

$$\varrho_n(\mathbf{r}') \prod_{i=1}^I \varphi\left(\frac{(m_{i,n} - m(r'_i))\sqrt{a\pi_i n}}{1}\right) \propto \exp(-n\psi(\mathbf{r}'; \mathbf{m}_n)),$$

where \propto denotes equality modulo normalization and

$$\psi(\mathbf{r}'; \mathbf{m}) := \frac{1}{2} \sum_{i=1}^I \left(\frac{\sigma(\mathbf{r}')}{\Delta} + a\pi_i (m(r'_i) - m_i)^2 \right).$$

The first inline equality follows from the specification of the prior ϱ_n in (10).

Since $m_{i,n} \rightarrow m(r_i)$ (a.s.), $\psi(\mathbf{r}'; \mathbf{m}_n)$ converges to $\psi(\mathbf{r}'; (m(r_i))_i)$, uniformly in \mathbf{r}' . Additionally, $\psi(\mathbf{r}'; (m(r_i))_i)$ as a function of \mathbf{r}' has the unique minimizer $\mathbf{q}^*(\mathbf{r})$ by assumption. Lemma 7 implies that the posterior formed given \mathbf{m}_n converges in probability to $\mathbf{q}^*(\mathbf{r})$. Since the support of the rewards is bounded, convergence in probability implies convergence in expected value, and thus the Bayesian estimate $\mathbb{E}[\mathbf{r} \mid \mathbf{m}_n]$ converges to $\mathbf{q}^*(\mathbf{r})$. \square

Proof of Proposition 7. By Proposition 6, the Bayesian estimate of \mathbf{r} converges to $\mathbf{q}^*(\mathbf{r})$. We write $\mathbf{q}^* = (q_i^*)_{i=1}^I$ as an abbreviation for $\mathbf{q}^*(\mathbf{r})$ and let $q^* = \sum_i p_i q_i^*$. The first-order condition applied to the minimization in (11) implies,

$$(q_i^* - q^*) + a\Delta(m(q_i^*) - m(r_i))m'(q_i^*) = 0, \tag{19}$$

for all $i = 1, \dots, I$, where we have used that $\pi_i = p_i$ and $\sum_i p_i (q_i^* - q^*) = q^* - q^* = 0$. We write σ^2 for $\sigma^2(\mathbf{r})$ and $\sigma^{*2} := \sum_{i=1}^I p_i (q_i^* - q^*)^2$ for the variance of \mathbf{q}^* . We will prove the following claims (see Footnote 19 for the definition of the $o(\cdot)$ convention):

Claim 1: Any function that is $o(r_i - r)$ or $o(q_i^* - r)$ is also $o(\sigma)$.

Claim 2: $q^* = r + o(\sigma)$.

Claim 3: $\sigma^{*2} = \frac{z(r)^2}{(1+z(r))^2} \sigma^2 + o(\sigma^2)$.

Claim 4: $q^* = r + \frac{1}{2} \frac{m''(r)}{m'(r)} \left(\sigma^2 + \left(\frac{2}{z(r)} - 1 \right) \sigma^{*2} \right) + o(\sigma^2)$.

To prove Claim 1, we provide a bound on the distance of r_i and r'_i from r . It follows from definition of σ^2 that $(r_i - r)^2 \leq \sigma^2/p_i$, and thus $|r_i - r| \leq \sigma/\sqrt{p_i}$. Therefore, any function that is $o(r_i - r)$ is also $o(\sigma)$. Bounding $|q_i^* - r|$ is complicated by the fact that \mathbf{q}^* is defined implicitly. We first establish a bound on $|q^* - r|$. Define \underline{m}' and \overline{m}' to be the minimum and the maximum of $m'(\cdot)$ on $[r, \bar{r}]$, respectively, and let $\underline{z} = a\Delta\underline{m}'^2$, $\bar{z} = a\Delta\overline{m}'^2$. We have $0 < \underline{m}' \leq \overline{m}' < +\infty$ and $0 < \underline{z} \leq \bar{z} < +\infty$ since $m'(\cdot)$ is continuous and strictly positive on the closed interval $[r, \bar{r}]$.

For fixed values of \mathbf{r} and \mathbf{q}^* define $z_i \in \mathbb{R}$ by

$$a\Delta m'(q_i^*) (m(q_i^*) - m_i(r_i)) = (q_i^* - r_i) z_i$$

whenever $q_i^* \neq r_i$, and $z_i := a\Delta m'^2(r_i)$ otherwise. It follows from its definition that $z_i \geq \underline{z}$ for all i . Then, equation (19) can be written as

$$0 = (q_i^* - q^*) + (q_i^* - r_i)z_i = (1 + z_i)(q_i^* - q^*) - (r_i - q^*)z_i,$$

and thus,

$$q_i^* - q^* = \frac{z_i}{1+z_i}(r_i - q^*) = \frac{z_i}{1+z_i}(r_i - r) + \frac{z_i}{1+z_i}(r - q^*). \quad (20)$$

Summing up the last equation weighted by p_i over i gives

$$0 = \sum_{i=1}^I \left(p_i \frac{z_i}{1+z_i}(r_i - r) \right) + (r - q^*) \sum_{i=1}^I \left(p_i \frac{z_i}{1+z_i} \right),$$

in which $0 < \frac{\underline{z}}{1+\underline{z}} \leq \frac{z_i}{1+z_i} < 1$. The triangle inequality implies

$$|q^* - r| \leq \frac{1+\underline{z}}{\underline{z}} \sum_{i=1}^I p_i |r_i - r| \leq \frac{1+\underline{z}}{\underline{z}} \sigma \sum_{i=1}^I \sqrt{p_i} \leq \frac{1+\underline{z}}{\underline{z}} I \sigma.$$

Returning to equation (20),

$$|q_i^* - r| \leq \frac{z_i}{1+z_i} |r_i - r| + \frac{z_i}{1+z_i} |r - q^*| + |q^* - r| < |r_i - r| + 2|r - q^*| \leq \left(p_i^{-1/2} + 2\frac{1+\underline{z}}{\underline{z}} I \right) \sigma.$$

We conclude that $|q_i^* - r| \leq \left(p_i^{-1/2} + 2\frac{1+\underline{z}}{\underline{z}}I\right)\sigma$ for any $\mathbf{r} \in [\underline{r}, \bar{r}]^I$, and thus any function that is $o(q_i^* - r)$ is also $o(\sigma)$. This establishes Claim 1.

We will prove the remaining claims by taking first- and second-order approximations of the first-order condition (19) for $\sigma > 0$ small. Since $m(\cdot)$ is twice differentiable, the functions m and m' can be expressed using first-order Taylor approximations around r :

$$\begin{aligned} m(r_i) &= m(r) + m'(r)(r_i - r) + o(\sigma), \\ m(q_i^*) &= m(r) + m'(r)(q_i^* - r) + o(\sigma), \\ m'(q_i^*) &= m'(r) + m''(r)(q_i^* - r) + o(\sigma), \end{aligned}$$

where we used Claim 1 to replace $o(r_i - r)$ and $o(q_i^* - r)$ by $o(\sigma)$. Equation (19) then implies

$$\begin{aligned} 0 &= (q_i^* - q^*) + a\Delta \left(m'(r)(q_i^* - r_i) + o(\sigma) \right) \left(m'(r) + m''(r)(q_i^* - r) + o(\sigma) \right) \\ &= (q_i^* - q^*) + a\Delta m'^2(r)(q_i^* - r_i) + o(\sigma), \end{aligned}$$

where we used that $(q_i^* - r_i)(q_i^* - r) = o(\sigma)$. The last inline equation can be written as

$$0 = (q_i^* - q^*) + z(r)(q_i^* - r_i) + o(\sigma). \quad (21)$$

Summing up these equations weighted by p_i , we get $0 = z(r)(q^* - r) + o(\sigma)$. Thus $|q^* - r| \leq \frac{1}{\underline{z}}o(\sigma)$, as needed for Claim 2.

We rewrite (21) as

$$(1 + z(r))(q_i^* - q^*) = z(r)(r_i - r) + z(r)(r - q^*) + o(\sigma) = z(r)(r_i - r) + o(\sigma),$$

where the second equality follows from Claim 2. Squaring both sides of the equation and summing up the equations weighted by p_i , we get

$$(1 + z(r))^2 \sigma^{*2} = z^2(r)\sigma^2 + o(\sigma^2),$$

where we used that $z(r) \leq \bar{z}$ and thus $z(r)(r_i - r)o(\sigma)$ is $o(\sigma^2)$. Claim 3 follows.

To prove Claim 4, we use the second-order Taylor approximation of $m(\cdot)$ around r :

$$\begin{aligned} m(q_i^*) &= m(r) + m'(r)(q_i^* - r) + \frac{1}{2}m''(r)(q_i^* - r)^2 + o(\sigma^2) \\ m(r_i) &= m(r) + m'(r)(r_i - r) + \frac{1}{2}m''(r)(r_i - r)^2 + o(\sigma^2). \end{aligned}$$

This implies the second-order approximation of the equation (19),

$$0 = (q_i^* - q^*) + a\Delta \left(m'(r)(q_i^* - r_i) + \frac{1}{2}m''(r)\left((q_i^* - r)^2 - (r_i - r)^2\right) + o(\sigma^2) \right) \\ \cdot \left(m'(r) + m''(r)(q_i^* - r) + o(\sigma) \right),$$

which we rewrite as

$$0 = (q_i^* - q^*) + z(r) \left((q_i^* - r_i) + \frac{1}{2} \frac{m''(r)}{m'(r)} \left((q_i^* - r)^2 - (r_i - r)^2 \right) \right) \left(1 + \frac{m''(r)}{m'(r)} (q_i^* - r) \right) + o(\sigma^2).$$

Summing up these equations weighted by p_i and dividing by $z(r)$, we arrive at

$$0 = (q^* - r) - \frac{1}{2} \frac{m''(r)}{m'(r)} \left(\sigma^2 - \sigma^{*2} + 2 \sum_{i=1}^I p_i (r_i - q_i^*) (q_i^* - r) \right) + o(\sigma^2). \quad (22)$$

Expressing $q_i^* - r_i$ from (21) allows us to write

$$\sum_{i=1}^I p_i (r_i - q_i^*) (q_i^* - r) = \frac{1}{z(r)} \sum_{i=1}^I p_i (q_i^* - r)^2 + o(\sigma^2) = \frac{1}{z(r)} \sigma^{*2} + o(\sigma^2),$$

where we used that $r = q^* + o(\sigma)$ for the second equality. Substituting the last inline equation back into (22) completes the proof of Claim 4.

Finally, substituting for σ^{*2} from Claim 3 into the expression from Claim 4 gives

$$q^* = r + \frac{1}{2} \frac{m''(r)}{m'(r)} \left(1 + \left(\frac{2}{z(r)} - 1 \right) \frac{z(r)^2}{(1+z(r))^2} \right) \sigma^2 + o(\sigma^2) \\ = r + \frac{1}{2} \frac{m''(r)}{m'(r)} \left(1 + \frac{2z(r) - z(r)^2}{(1+z(r))^2} \right) \sigma^2 + o(\sigma^2),$$

and using $1 + \frac{2z(r) - z(r)^2}{(1+z(r))^2} = \frac{1+4z(r)}{(1+z(r))^2}$, we obtain (12), concluding the proof. \square